

ABIRATERONE

Products Affected

- *abiraterone acetate*

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Castration-resistant metastatic prostate cancer and used in combination with prednisone, or B.) High risk, castration-sensitive metastatic prostate cancer and used in combination with prednisone |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or urologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ACITRETIN

Products Affected

- *acitretin*

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Severely impaired liver or kidney function, B.) Chronic abnormally elevated blood lipid values, C.) Concomitant use of methotrexate or tetracyclines, D.) Pregnancy |
| Required Medical Information | Diagnosis of severe, recalcitrant psoriasis (including plaque, guttate, erythrodermic palmar- plantar and pustular) AND patient must have tried and failed, contraindication or intolerance to one formulary first line agent (e.g., Topical Corticosteroids (betamethasone, fluocinonide, desoximetasone), Topical Calcipotriene/Calcitriol, Topical Calcipotriene, OR Topical Tazarotene) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a dermatologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ACTIMMUNE

Products Affected

- ACTIMMUNE

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Chronic granulomatous disease for use in reducing the frequency and severity of serious infections, or B.) Severe, malignant osteopetrosis (SMO) |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | B vs D determination required per CMS guidance |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ADEMPAS

Products Affected

- ADEMPAS

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Concomitant administration with nitrates or nitric oxide donors (such as amyl nitrate) in any form, B.) Concomitant administration with phosphodiesterase inhibitors, including specific PDE-5 inhibitors (such as sildenafil, tadalafil, or vardenafil) or non-specific PDE inhibitors (such as dipyridamole or theophylline), C.) Pregnancy, or D.) Patients with pulmonary hypertension associated with idiopathic interstitial pneumonia |
| Required Medical Information | Diagnosis of one of the following A.) Pulmonary arterial hypertension (WHO group I) and diagnosis was confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.), or B.) Chronic thromboembolic pulmonary hypertension (CTEPH, WHO group 4) and patient has persistent or recurrent disease after surgical treatment (e.g., pulmonary endarterectomy) or has CTEPH that is inoperable (Female patients must be enrolled in the ADEMPAS REMS program) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a cardiologist or pulmonologist |
| Coverage Duration | Initial: 6 months, Renewal: 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ALECENSA

Products Affected

- ALECENSA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of metastatic anaplastic lymphoma kinase (ALK) positive non-small cell lung cancer as detected by an FDA-approved test |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ALPHA-1 PROTEINASE INHIBITOR

Products Affected

- PROLASTIN-C INTRAVENOUS
SOLUTION RECONSTITUTED

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Immunoglobulin A (IgA) deficiency with antibodies against IgA |
| Required Medical Information | Diagnosis of alpha-1 proteinase inhibitor (alpha-1 antitrypsin) deficiency in adult patients with emphysema |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a pulmonologist |
| Coverage Duration | 12 months |
| Other Criteria | B vs D determination required per CMS guidance |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ALUNBRIG

Products Affected

- ALUNBRIG

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of anaplastic lymphoma kinase-positive (ALK) metastatic non-small cell lung cancer (NSCLC) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

AMBRISENTAN

Products Affected

- ambrisentan*

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Pregnancy, or B.) Idiopathic pulmonary fibrosis (IPF), including those with pulmonary hypertension |
| Required Medical Information | Diagnosis of pulmonary arterial hypertension classified as WHO Group I, confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a cardiologist or pulmonologist |
| Coverage Duration | Initial: 6 months, Renewal: 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ARCALYST

Products Affected

- ARCALYST

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Cryopyrin-associated periodic syndromes (CAPS), including familial cold autoinflammatory syndrome (FCAS) and Muckle-Wells Syndrome (MWS), B.) Deficiency of interleukin-1 receptor antagonist (DIRA) and patient requires maintenance therapy for remission, or C.) Recurrent pericarditis (RP) and reduction in risk of recurrence |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | B vs D determination required per CMS guidance |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ARIKAYCE

Products Affected

- ARIKAYCE

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | Known sensitivity to any aminoglycoside |
| Required Medical Information | Diagnosis of pulmonary Mycobacterium avium complex (MAC) infection and used as part of a combination antibacterial regimen in treatment refractory patients (greater than 6 months of a multidrug background regimen) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an infectious disease specialist or pulmonologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

AURYXIA

Products Affected

- AURYXIA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | Iron overload syndrome (e.g. hemochromatosis) |
| Required Medical Information | Diagnosis of hyperphosphatemia in patients with chronic kidney disease (CKD) on dialysis |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist or nephrologist |
| Coverage Duration | 12 months |
| Other Criteria | Ferric Citrate is NOT approvable for iron deficiency anemia per Part D law |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

AUSTEDO

Products Affected

- AUSTEDO

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Suicidal ideation and/or untreated or inadequately treated depression in a patient with Huntington's Disease, B.) Hepatic impairment, C.) Concomitant use of MAOIs, reserpine, tetrabenazine, or valbenazine |
| Required Medical Information | Diagnosis of one of the following A.) Chorea associated with Huntington's disease (Huntington's chorea), or B.) Tardive dyskinesia |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a neurologist or psychiatrist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

AYVAKIT

Products Affected

- AYVAKIT

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Unresectable or metastatic gastrointestinal stromal tumor, with a platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation, including PDGFRA D842V mutations, or B.) Advanced Systemic Mastocytosis (AdvSM), including aggressive systemic mastocytosis (ASM), systemic mastocytosis with an associated hematological neoplasm (SMAHN), or mast cell leukemia (MCL), and platelet count of at least 50,000/mcL |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

BALVERSA

Products Affected

- BALVERSA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of locally advanced or metastatic urothelial carcinoma and both of the following 1.) Susceptible fibroblast growth factor receptor (FGFR)3 or FGFR2 genetic alterations confirmed by an FDA-approved diagnostic test, and 2.) Patient has progressed during or following at least one line of prior platinum-containing chemotherapy, including within 12 months of neoadjuvant or adjuvant platinum-containing chemotherapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or urologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

BENLYSTA

Products Affected

- BENLYSTA SUBCUTANEOUS

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Active, autoantibody-positive, system lupus erythematosus (SLE), or B.) Active lupus nephritis and patient is receiving standard therapy |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with a nephrologist or rheumatologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

BESREMI

Products Affected

- BESREMI

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Existence of, or history of severe psychiatric disorders (severe depression, suicidal ideation, or suicide attempt), B.) Hypersensitivity to interferons including interferon alfa-2b or excipients, C.) Hepatic impairment (Child-Pugh B or C), D.) History or presence of active serious or untreated autoimmune disease, or E.) Immunosuppressed transplant recipients |
| Required Medical Information | Diagnosis of polycythemia vera |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

BEXAROTENE GEL

Products Affected

- *bexarotene external*

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of primary cutaneous T-cell lymphoma (CTCL Stage 1A/1B) and patient had an inadequate response, is intolerant to, or has a contraindication to at least one prior systemic therapy (e.g., corticosteroids) indicated for cutaneous manifestations of CTCL |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or dermatologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

BEXAROTENE ORAL

Products Affected

- bexarotene oral*

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | Pregnancy |
| Required Medical Information | Diagnosis of cutaneous T-cell lymphoma (CTCL) and patient is not a candidate for or had an inadequate response, is intolerant to, or has a contraindication to at least one prior systemic therapy (e.g., corticosteroids) for cutaneous manifestations of CTCL |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with an dermatologist, oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

BOSENTAN

Products Affected

- bosentan*

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Concomitant cyclosporine A or glyburide therapy, or B.) Pregnancy |
| Required Medical Information | Diagnosis of pulmonary arterial hypertension (WHO Group I) and patient has New York Heart Association (NYHA) Functional Class II-IV, confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e. g., patient is frail, elderly, etc.) |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with a cardiologist or pulmonologist |
| Coverage Duration | Initial: 6 months, Renewal: 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

BOSULIF

Products Affected

- BOSULIF

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Chronic, accelerated, or blast phase Philadelphia chromosome-positive (Ph+) chronic myelogenous leukemia (CML) with resistance or inadequate response to prior therapy, or B.) Newly diagnosed chronic phase Philadelphia chromosome-positive (Ph+) CML |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

BRAFTOVI

Products Affected

- BRAFTOVI ORAL CAPSULE 75 MG

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) unresectable or metastatic melanoma with documented BRAF V600E or V600K mutation as detected by a FDA-approved test and used in combination with binimetinib, or B.) metastatic colorectal cancer with documented BRAF V600E mutation as detected by a FDA-approved test, patient has received prior therapy, and braftovi used in combination with cetuximab. |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

BRUKINSA

Products Affected

- BRUKINSA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following: A.) mantle cell lymphoma (MCL) and patient has received at least one prior therapy or B.) Treatment of adult patients with Waldenstrom macroglobulinemia or C.) Treatment of adult patients with relapsed or refractory marginal zone lymphoma who have received at least one anti-CD20-based regimen |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

BYLVAY

Products Affected

- BYLVAY
- BYLVAY (PELLETS) ORAL CAPSULE SPRINKLE 200 MCG

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of progressive familial intrahepatic cholestasis-associated pruritus |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

CABLIVI

Products Affected

- CABLIVI

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of acquired thrombotic thrombocytopenic purpura (aTTP) and used in combination with plasma exchange and immunosuppression therapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist or oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

CABOMETYX

Products Affected

- CABOMETYX

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Advanced renal cell carcinoma, B.) Advanced hepatocellular carcinoma (HCC) and patient has been previously treated with sorafenib, C.) Advanced renal cell carcinoma and used as first line treatment in combination with nivolumab or D.) treatment of adults and pediatric patients 12 years and older with locally advanced or metastatic differentiated thyroid cancer that has progressed following VEGFR-targeted therapy and who are radioactive iodine-refractory or ineligible |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

CALQUENCE

Products Affected

- CALQUENCE

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Mantle cell lymphoma (MCL) and patient has received at least 1 prior therapy, B.) Chronic lymphocytic leukemia (CLL), or C.) Small lymphocytic lymphoma (SLL) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

CAMZYOS

Products Affected

- CAMZYOS

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of symptomatic New York Heart Association (NYHA) class II-III obstructive hypertrophic cardiomyopathy (HCM) in adult patients |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

CAPRELSA

Products Affected

- CAPRELSA

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | Congenital long QT syndrome |
| Required Medical Information | Diagnosis of metastatic or unresectable locally advanced medullary thyroid cancer (MTC) AND disease is symptomatic or progressive |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

CARGLUMIC ACID

Products Affected

- carglumic acid oral tablet soluble*

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) N-acetyl glutamate synthase (NAGS) deficiency (confirmed by appropriate genetic testing) with acute or chronic hyperammonemia, or B.) Propionic or methylmalonic acidemia with acute hyperammonemia |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

CAYSTON

Products Affected

- CAYSTON

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of cystic fibrosis (confirmed by appropriate diagnostic or genetic testing) and patient has Pseudomonas aeruginosa lung infection confirmed by positive culture |
| Age Restrictions | 7 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

CNS STIMULANTS

Products Affected

- armodafinil*
- modafinil*

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Obstructive sleep apnea (OSA) confirmed by sleep lab evaluation, B.) Narcolepsy confirmed by sleep lab evaluation, or C.) Shift work disorder (SWD) |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

COMETRIQ

Products Affected

- COMETRIQ (100 MG DAILY DOSE)
ORAL KIT 80 & 20 MG
- COMETRIQ (140 MG DAILY DOSE)
ORAL KIT 3 X 20 MG & 80 MG
- COMETRIQ (60 MG DAILY DOSE)

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of progressive, metastatic medullary thyroid cancer |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

COPIKTRA

Products Affected

- COPIKTRA

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Relapsed or refractory chronic lymphocytic leukemia (CLL), or B.) Relapsed or refractory small lymphocytic lymphoma (SLL). For CLL, or SLL, the patient must have history of at least 2 prior therapies |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

CORLANOR

Products Affected

- CORLANOR ORAL TABLET

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Decompensated acute heart failure, B.) hypotension (i.e. blood pressure less than 90/50 mmHg), C.) sick sinus syndrome or sinoatrial block or 3rd degree AV block (unless a functioning demand pacemaker is present), D.) bradycardia (i.e., resting heart rate less than 60 bpm prior to treatment), E.) Severe hepatic impairment (Child-Pugh C), F.) Pacemaker dependent (heart rate maintained exclusively by the pacemaker), G.) Concomitant use of strong CYP3A4 inhibitors |
| Required Medical Information | Diagnosis of one of the following A.) stable, symptomatic chronic heart failure with left ventricular ejection fraction 35% or less, who are in sinus rhythm with resting heart rate 70 beats per minute or more and either are on maximally tolerated doses of beta-blockers or have a contraindication to beta-blocker use, or B.) stable, symptomatic heart failure due to dilated cardiomyopathy in patients who are in sinus rhythm with an elevated heart rate |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

COSENTYX

Products Affected

- COSENTYX (300 MG DOSE)
- COSENTYX SENSOREADY (300 MG)
- COSENTYX SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 75 MG/0.5ML

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Ankylosing spondylitis and patient has trial and failure, contraindication, or intolerance to two preferred products, (i.e. Humira, Enbrel), B.) Moderate to severe plaque psoriasis in adults and patient has trial and failure, contraindication, or intolerance to two preferred products, (i.e. Humira, Enbrel, Skyrizi, Stelara), C.) Moderate to severe plaque psoriasis in patients 6 years to less than 18 years of age and patient has trial and failure, contraindication, or intolerance to two preferred products, (i.e. Enbrel, Stelara), D.) Active psoriatic arthritis in adult patient and has trial and failure, contraindication, or intolerance to two preferred products, (i.e. Humira, Enbrel, Skyrizi, Stelara), E.) Active psoriatic arthritis in patients 2 years to less than 18 years of age, F.) Non-radiographic axial spondyloarthritis or G.) Active enthesitis-related arthritis |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | Screening for latent tuberculosis infection is required prior to initiation of treatment |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

COTELLIC

Products Affected

- COTELLIC

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.)unresectable or metastatic malignant melanoma with BRAF V600E OR V600K mutation, and documentation of combination therapy with vemurafenib (Zelboraf), or B.) Histiocytic neoplasms |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

CYSTAGON

Products Affected

- CYSTAGON

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | Known serious hypersensitivity to penicillamine or cysteamine |
| Required Medical Information | Diagnosis of nephropathic cystinosis confirmed by the presence of increased cystine concentration in leukocytes or by genetic testing |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

CYSTEAMINE OPHTH

Products Affected

- CYSTADROPS
- CYSTARAN

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of cystinosis and patient has corneal cystine crystal accumulation |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

DALFAMPRIDINE

Products Affected

- dalfampridine er*

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) History of seizure. B.) Moderate or severe renal impairment (creatinine clearance less than or equal to 50 mL/minute) |
| Required Medical Information | Diagnosis of multiple sclerosis and patient must demonstrate sustained walking impairment, but with the ability to walk 25 feet (with or without assistance) prior to starting dalfampridine |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a neurologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

DAURISMO

Products Affected

- DAURISMO

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of newly diagnosed acute myeloid leukemia (AML) and used in combination with cytarabine in patients 75 years of age or older OR in patients that have comorbidities that preclude use of intensive induction chemotherapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

DEFERASIROX

Products Affected

- *deferasirox granules*
- *deferasirox oral tablet*
- *deferasirox oral tablet soluble*

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Creatinine clearance less than 40 mL/min, B.) Poor performance status, C.) Platelet count less than 50 x 10 ⁹ /L, D.) Advanced malignancy, E.) High-risk myelodysplastic syndrome (MDS) |
| Required Medical Information | Diagnosis of one of the following A.) Chronic iron overload in patients with non-transfusion-dependent thalassemia syndromes who have liver iron concentrations of at least 5 mg Fe/g dry weight AND serum ferritin level greater than 300 mcg/L, or B.) Chronic iron overload due to blood transfusions (transfusion hemosiderosis) as evidenced by transfusion of at least 100 mL/kg packed red blood cells AND serum ferritin level greater than 1000 mcg/L |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

DEFERIPRONE

Products Affected

- *deferiprone*
- FERRIPROX TWICE-A-DAY
- FERRIPROX ORAL SOLUTION

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Must meet all of the following 1.) Diagnosis of transfusional iron overload due to thalassemia syndromes, sickle cell disease, or other anemias, 2.) Patient has failed prior chelation therapy, and 3.) Patient has an absolute neutrophil count greater than $1.5 \times 10^9/L$ |
| Age Restrictions | 3 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

DIACOMIT

Products Affected

- DIACOMIT

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of severe myoclonic epilepsy in infancy (Dravet syndrome) in patients taking clobazam |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with a neurologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

DICLOFENAC TOPICAL

Products Affected

- diclofenac sodium external gel 3 %*

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------------|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of Actinic keratosis |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

DIFICID

Products Affected

- DIFICID

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of diarrhea associated with clostridioides difficile infection and patient has had an inadequate treatment response, intolerance, or contraindication to generic oral vancomycin |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 4 weeks |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

DIMETHYL FUMARATE

Products Affected

- dimethyl fumarate oral*
- dimethyl fumarate starter pack*

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a neurologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

DOJOLVI

Products Affected

- DOJOLVI

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of Long-chain fatty acid oxidation disorder (LC-FAOD) |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

DRONABINOL

Products Affected

- dronabinol*

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | Sesame oil hypersensitivity |
| Required Medical Information | Diagnosis of one of the following A.) Anorexia associated to AIDS, or B.) Chemotherapy-induced nausea and vomiting |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 6 months |
| Other Criteria | B vs D determination required per CMS guidance |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

DROXIDOPA

Products Affected

- droxidopa*

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of symptomatic neurogenic orthostatic hypotension (nOH) caused by primary autonomic failure (e.g., Parkinson disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, or non-diabetic autonomic neuropathy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

DUPIXENT

Products Affected

- DUPIXENT

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Moderate to severe atopic dermatitis and if patient is 2 years or older has trial/failure, contraindication, or intolerance to two of the following 1.) Topical corticosteroid, and/or 2.) Topical calcineurin inhibitor, B.) Eosinophilic phenotype or oral corticosteroid- dependent moderate to severe asthma and used as an adjunct treatment, or C.) Chronic rhinosinusitis with nasal polyposis and used as an adjunct treatment, D.) Eosinophilic esophagitis, or E.) Prurigo nodularis |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

EMGALITY

Products Affected

- EMGALITY

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Chronic or episodic migraine disorder and patient has documented trial, inadequate response, or contraindication to at least 2 generic formulary drugs used for migraine prevention (i.e., propranolol, topiramate, divalproex, timolol), or B.) Episodic cluster headache |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ENBREL

Products Affected

- ENBREL MINI
- ENBREL SUBCUTANEOUS SOLUTION 25 MG/0.5ML
- ENBREL SUBCUTANEOUS SOLUTION PREFILLED SYRINGE
- ENBREL SURECLICK SUBCUTANEOUS SOLUTION AUTO-INJECTOR

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Moderate to severe polyarticular juvenile idiopathic arthritis, C.) Psoriatic arthritis, D.) Ankylosing spondylitis, or E.) Moderate to severe chronic plaque psoriasis in patients who are candidates for systemic therapy or phototherapy |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | Screening for latent tuberculosis infection is required prior to initiation of treatment |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ENDARI

Products Affected

- ENDARI

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of sickle cell disease AND one of the following 1.) Patient has acute complications and is being treated with Hydroxyurea, or 2.) Patient has acute complications and is unable to tolerate Hydroxyurea |
| Age Restrictions | 5 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ENSPRYNG

Products Affected

- ENSPRYNG

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Active Hepatitis B infection, or B.) Active or untreated latent tuberculosis |
| Required Medical Information | Diagnosis of neuromyelitis optica spectrum disorder (NMOSD) in patients who are anti-aquaporin-4 (AQP4) antibody positive |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a neurologist, immunologist, or ophthalmologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

EPIDIOLEX

Products Affected

- EPIDIOLEX

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Lennox-Gastaut syndrome, B.) Severe myoclonic epilepsy in infancy (Dravet syndrome), or C.) Seizures associated with tuberous sclerosis complex (TSC) |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with a neurologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

EPOETIN THERAPY

Products Affected

- RETACRIT INJECTION SOLUTION UNIT/ML, 4000 UNIT/ML, 40000
10000 UNIT/ML, 10000 UNIT/ML(1ML), UNIT/ML
2000 UNIT/ML, 20000 UNIT/ML, 3000

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Non-myeloid neoplastic disease and utilized for the treatment of chemotherapy induced anemia, B.) HIV infection and utilized for the treatment of zidovudine induced anemia, C.) Chronic kidney disease resulting in anemia, or D.) High risk surgical candidate status at risk for perioperative blood loss and undergoing elective, noncardiac, or nonvascular surgery |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | B vs D determination required per CMS guidance |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ERIVEDGE

Products Affected

- ERIVEDGE

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | Pregnancy |
| Required Medical Information | Diagnosis of one of the following A.) Metastatic basal cell carcinoma, or B.) Locally advanced basal cell carcinoma that has recurred following surgery or the patient is not a candidate for surgery or radiation |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ERLEADA

Products Affected

- ERLEADA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Nonmetastatic, castration-resistant prostate cancer, or B.) Metastatic, castration-sensitive prostate cancer |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or urologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ERLOTINIB

Products Affected

- *erlotinib hcl*

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Locally advanced, unresectable, or metastatic pancreatic cancer and erlotinib will be used in combination with gemcitabine, B.) Locally advanced or metastatic non-small cell lung cancer with epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility AND one of the following 1.) Erlotinib will be used as first-line treatment, 2.) Failure with at least one prior chemotherapy regimen, or 3.) No evidence of disease progression after four cycles of first-line platinum-based chemotherapy and erlotinib will be used as maintenance treatment |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ESBRIET

Products Affected

- ESBRIET ORAL CAPSULE

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of idiopathic pulmonary fibrosis |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a pulmonologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

EVEROLIMUS

Products Affected

- *everolimus oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg*

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Hypersensitivity to everolimus or excipients, or B.) Hypersensitivity to rapamycin derivatives (e.g. sirolimus) |
| Required Medical Information | Diagnosis of one of the following A.) Renal angiomyolipoma and tuberous sclerosis complex (TSC) not requiring immediate surgery, B.) Advanced hormone receptor-positive, HER2 negative breast cancer in postmenopausal women and taken in combination with exemestane, after failure with letrozole or anastrozole, C.) Progressive, well-differentiated, nonfunctional neuroendocrine tumors of gastrointestinal or lung origin and disease is unresectable, locally advanced, or metastatic, D.) Pancreatic progressive neuroendocrine tumors and disease is unresectable, locally advanced, or metastatic, E.) Advanced renal cell carcinoma (RCC) after failure with sunitinib or sorafenib, F.) Subependymal giant cell astrocytoma (SEGA) associated with tuberous sclerosis complex in patients who are not candidates for curative surgical resection |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or neurologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

EVEROLIMUS SUSPENSION

Products Affected

- *everolimus oral tablet soluble*

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Hypersensitivity to everolimus , or B.) Hypersensitivity to rapamycin derivatives (e.g. sirolimus) |
| Required Medical Information | Diagnosis of one of the following A.) Tuberous sclerosis complex (TSC)-associated partial-onset seizures, or B.) Subependymal giant cell astrocytoma (SEGA) associated with tuberous sclerosis complex in patients who are not candidates for curative surgical resection |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or neurologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

EVRYSDI

Products Affected

- EVRYSDI

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of spinal muscular atrophy (SMA) |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with a neurologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

EXKIVITY

Products Affected

- EXKIVITY

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of locally advanced or metastatic non-small cell lung cancer (NSCLC) with EGFR exon 20 insertion mutations (as confirmed by an FDA-approved test) AND whose disease has progressed on or after platinum-based chemotherapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

FEBUXOSTAT

Products Affected

- *febuxostat*

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Concomitant use of azathioprine or mercaptopurine |
| Required Medical Information | Diagnosis of Gout and all of the following 1.) documented inadequate treatment response, adverse event, or contraindication to maximally titrated dose of Allopurinol, and 2.) patients with established cardiovascular disease, prescriber attests that benefit of treatment outweighs the risk of treatment |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

FENTANYL ORAL

Products Affected

- fentanyl citrate buccal lozenge on a handle*

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Management of acute or postoperative pain (including headache/migraine, dental pain, and use in the emergency room), B.) Use in opioid non-tolerant patients, C.) Known or suspected gastrointestinal obstruction, including paralytic ileus, D.) Acute or severe bronchial asthma and used in an unmonitored setting (absence of resuscitative equipment) |
| Required Medical Information | Must meet all of the following 1.) Diagnosis of cancer-related breakthrough pain, 2.) Patient is currently receiving/tolerant to around-the-clock opioid therapy for persistent cancer pain, and 3.) Patient and prescriber are enrolled in the TIRF REMS Access Program |
| Age Restrictions | 16 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

FENTANYL PATCH

Products Affected

- *fentanyl*

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Management of acute or postoperative pain (including headache/migraine, dental pain, and use in the emergency room), B.) Mild or intermittent pain management, C.) Use in opioid non-tolerant patients, D.) Known or suspected gastrointestinal obstruction, including paralytic ileus, E.) Acute or severe bronchial asthma and used in an unmonitored setting (absence of resuscitative equipment) |
| Required Medical Information | Must meet all of the following 1.) Patient is opioid tolerant (taking for one week or longer at least 60mg of morphine or equivalent daily) and 2.) Patient has tried at least one extended release oral opioids or is unable to take extended release oral opioids secondary to allergy, adverse events, swallowing difficulty, or uncontrollable nausea/vomiting |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

FINGOLIMOD

Products Affected

- fingolimod hcl*

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Recent (within the last 6 months) occurrence of: myocardial infarction, unstable angina, stroke, transient ischemic attack, decompensated heart failure requiring hospitalization, or Class III/IV heart failure, B.) History or presence of Mobitz Type II 2nd degree or 3rd degree AV block or sick sinus syndrome, unless patient has a pacemaker, C.) Baseline QTC interval greater than or equal to 500 milliseconds, D.) Receiving concurrent treatment with Class Ia or Class III anti-arrhythmic drugs (quinidine, procainamide, amiodarone, sotalol) |
| Required Medical Information | Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis |
| Age Restrictions | 10 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a neurologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

FINTEPLA

Products Affected

- FINTEPLA

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Concomitant use of an MAOI, or B.) Use within 14 days of discontinuing an MAOI |
| Required Medical Information | Diagnosis of Severe myoclonic epilepsy in infancy (Dravet syndrome) or seizures associated with Lennox-Gastaut syndrome |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with a neurologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

FOTIVDA

Products Affected

- FOTIVDA

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of relapsed or refractory advanced renal cell cancer (RCC) following 2 or more prior systemic therapies |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

GALAFOLD

Products Affected

- GALAFOLD

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of Fabry disease with an amenable galactosidase alpha gene (GLA) mutation |
| Age Restrictions | 16 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

GATTEX

Products Affected

- GATTEX

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of short bowel syndrome and patient is dependent on parenteral support |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

GAVRETO

Products Affected

- GAVRETO

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Metastatic RET fusion-positive non-small cell lung cancer (NSCLC) as detected by an FDA approved test, B.) Advanced or metastatic RET-mutant medullary thyroid cancer and patient requires systemic therapy, or C.) Advanced or metastatic RET fusion-positive thyroid and patient requires systemic therapy and is radioactive iodine-refractory, when radioactive iodine is appropriate |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

GILOTRIF

Products Affected

- GILOTRIF

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Metastatic non-small cell lung cancer (NSCLC) in patients whose tumors have non-resistant epidermal growth factor receptor (EGFR) mutations as detected by an FDA-approved test, or B.) Metastatic squamous NSCLC with progression after platinum-based chemotherapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

GLATIRAMER

Products Affected

- COPAXONE SUBCUTANEOUS
SOLUTION PREFILLED SYRINGE

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a neurologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

GLEOSTINE

Products Affected

- GLEOSTINE ORAL CAPSULE 10 MG, 100 MG, 40 MG

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Must meet one of the following: A.) Hodgkin's disease in patient who has relapsed during or failed to respond to primary therapy and is being used in combination with other agents OR B.) Intracranial tumor |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

GROWTH HORMONE

Products Affected

- OMNITROPE SUBCUTANEOUS SOLUTION CARTRIDGE
- OMNITROPE SUBCUTANEOUS SOLUTION RECONSTITUTED

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Use for growth promotion in pediatric patients with closed epiphyses, B.) Acute critical illness caused by complications following open-heart or abdominal surgery, multiple accidental trauma, or acute respiratory failure, C.) Active malignancy, D.) Active proliferative or severe nonproliferative diabetic retinopathy, E.) Prader-Willi Syndrome in patients who are severely obese, have a history of upper airway obstruction or sleep apnea, or have severe respiratory impairment |
| Required Medical Information | Diagnosis of pediatric indication: A.) GHD and bone age at least 1 year or 2 standard deviations (SD) delayed compared with chronological age and 2 stim tests with peak GH secretion below 10 ng/mL or IGF-1/IGFBP3 level more than 2 SDS below mean if CNS pathology, h/o irradiation, or proven genetic cause, B.) SGA and birth weight or length 2 or more SDS below mean for gestational age and fails to manifest catch up growth by age 2 (height 2 or more SDS below mean for age and gender), C.) CRI and metabolic abnormalities have been corrected, and patient has not had renal transplant D.) SHOX deficiency or Noonan syndrome E.) PWS confirmed by genetic testing, F.) Turner Syndrome confirmed by chromosome analysis. For GHD, CRI, SHOX deficiency, Noonan syndrome, and PWS one of the following height more than 3 SDS below mean for age and gender, or height more than 2 SDS below mean with GV more than 1 SDS below mean, or GV over 1 year 2 SDS below mean. OR Diagnosis of an adult indication: A.) childhood- or adult-onset GHD confirmed by 2 standard GH stim tests (provide assay): 1 test must be insulin tolerance test (ITT) with blood glucose nadir less than 40 mg/dL (2.2 mmol/L). If contraindicated, use a standardized stim test (i.e. arginine plus GH releasing hormone [preferred], glucagon, arginine), B.) GHD with at least 1 other pituitary hormone deficiency and failed at least 1 GH stim test (ITT preferred), C.) GHD with panhypopituitarism (3 or more pituitary hormone deficiencies), D.) GHD with irreversible hypothalamic-pituitary structural lesions due to tumors, surgery or radiation of pituitary or hypothalamus region AND a subnormal IGF-1 (after at least 1 month off GH therapy) AND Objective evidence of GHD complications, such as: low bone density, increased visceral fat mass, or cardiovascular complications AND |

| PA Criteria | Criteria Details |
|-------------------------|---|
| | Completed linear growth (GV less than 2 cm/year) AND GH has been discontinued for at least 1 month (if previously receiving GH) |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with an Endocrinologist or Nephrologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

HEPATITIS C

Products Affected

- MAVYRET
- *sofosbuvir-velpatasvir*
- VOSEVI

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of HCV genotype, subtype and quantitative HCV RNA (viral load) testing any time prior to therapy. Must document cirrhosis status, prior treatment history (if any), and planned duration of treatment. All genotypes will require trial/failure, contraindication to, or intolerance to Mavyret or Sofosbuvir-Velpatasvir prior to the approval of Vosevi. |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with a gastroenterologist, hepatologist, or infectious disease specialist |
| Coverage Duration | Duration of approval per AASLD Guidelines |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

HUMIRA

Products Affected

- HUMIRA PEDIATRIC CROHNS START SUBCUTANEOUS PREFILLED SYRINGE KIT 80 MG/0.8ML, 80 MG/0.8ML & 40MG/0.4ML
- HUMIRA PEN SUBCUTANEOUS PEN-INJECTOR KIT
- HUMIRA PEN-CD/UC/HS STARTER
- HUMIRA PEN-PEDIATRIC UC START
- HUMIRA PEN-PS/UV/ADOL HS START SUBCUTANEOUS PEN-INJECTOR KIT 40 MG/0.8ML
- HUMIRA PEN-PSOR/UEIT STARTER
- HUMIRA SUBCUTANEOUS PREFILLED SYRINGE KIT 10 MG/0.1ML, 20 MG/0.2ML, 40 MG/0.4ML, 40 MG/0.8ML

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Moderate to severe polyarticular juvenile idiopathic arthritis, C.) Psoriatic arthritis, D.) Ankylosing spondylitis, E.) Moderate to severe chronic plaque psoriasis in patients who are candidates for systemic therapy or phototherapy and when other systemic therapies are medically less appropriate, F.) Moderate to severe Crohn's disease in patients who have had an inadequate response to conventional therapy, G.) Moderate to severe ulcerative colitis in patients who have had an inadequate response to immunosuppressants (e.g. corticosteroids, azathioprine), H.) Non-infectious uveitis (including intermediate, posterior, and panuveitis), or I.) Moderate to severe hidradenitis suppurativa |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | Screening for latent tuberculosis infection is required prior to initiation of treatment |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |

| PA Criteria | Criteria Details |
|------------------------|------------------|
| Part B Prerequisite | No |

HYFTOR

Products Affected

- HYFTOR

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of Facial angiofibroma associated with tuberous sclerosis |
| Age Restrictions | 6 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

IBRANCE

Products Affected

- IBRANCE

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer used in combination with fulvestrant and disease has progressed following endocrine therapy, or B.) Advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer and used in combination with an aromatase inhibitor in a male or postmenopausal female patient as initial endocrine-based therapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ICATIBANT

Products Affected

- FIRAZYR
- icatibant acetate*

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following and used as treatment for acute attacks A.) Hereditary angioedema (HAE) with C1 inhibitor deficiency (Type 1) confirmed by laboratory testing, or B.) HAE with C1 inhibitor dysfunction (Type 2) confirmed by laboratory testing, or C.) HAE with normal C1 inhibitor (Type 3) confirmed by laboratory testing and one of the following 1.) Positive test for an F12, angiopoietin-1, or plasminogen gene mutation, or 2.) Family history of angioedema and the angioedema was refractory to a trial of an antihistamine for at least one month |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an allergist, hematologist, or immunologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ICLUSIG

Products Affected

- ICLUSIG

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Chronic phase, accelerated phase, or blast phase chronic myeloid leukemia (CML) in adult patients who are T315I-positive or for whom no other tyrosine kinase inhibitor therapy is indicated, B.) Chronic phase, chronic myeloid leukemia (CML) in adult patients with resistance or intolerance to at least two prior kinase inhibitors, or C.) Philadelphia chromosome-positive acute lymphoblastic leukemia (Ph+ ALL) in adult patients who are T315I-positive or for whom no other tyrosine kinase inhibitor therapy is indicated |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

IDHIFA

Products Affected

- IDHIFA

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of relapsed or refractory acute myeloid leukemia (AML) with an isocitrate dehydrogenase 2 (IDH2) mutation as detected by an FDA approved test |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

IMATINIB

Products Affected

- imatinib mesylate*

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Philadelphia chromosome-positive chronic myelogenous leukemia (Ph+ CML), B.) Ph+ acute lymphoblastic leukemia (ALL), C.) Gastrointestinal stromal tumor (GIST) where patient has documented c-KIT (CD117) positive unresectable or metastatic malignant GIST or patient had resection of c-KIT positive GIST and imatinib will be used as an adjuvant therapy, D.) Dermatofibrosarcoma protuberans that is unresectable, recurrent, or metastatic, E.) Hypereosinophilic syndrome or chronic eosinophilic leukemia, F.) Myelodysplastic syndrome or myeloproliferative disease associated with platelet-derived growth factor receptor gene re-arrangements, or G.) Aggressive systemic mastocytosis without the D816V c-KIT mutation or with c-KIT mutational status unknown |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

IMBRUVICA

Products Affected

- IMBRUVICA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Mantle cell lymphoma (MCL) and patient has received at least one prior therapy, B.) Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL), C.) Chronic lymphocytic leukemia (CLL)/Small lymphocytic lymphoma (SLL) with 17p deletion, D.) Waldenstrom's macroglobulinemia (WM), E.) Marginal zone lymphoma (MZL) and patient requires systemic therapy and has received at least one prior anti-CD20-based therapy, or F.) Chronic graft vs host disease (cGVHD) after failure of at least one first-line corticosteroid therapy |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

INCRELEX

Products Affected

- INCRELEX

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Active or suspected malignancy, B.) Use for growth promotion in patients with closed epiphyses, or C.) Intravenous administration |
| Required Medical Information | Diagnosis of one of the following A.) Severe primary insulin-like growth factor-1 (IGF-1) deficiency and utilized for pediatric treatment of growth failure, or B.) Growth hormone (GH) gene deletion and patient has developed neutralizing antibodies to GH and utilized for pediatric treatment of growth failure |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

INLYTA

Products Affected

- INLYTA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Advanced renal cell carcinoma and patient failed one or more systemic therapies for renal cell carcinoma (e.g., sunitinib-, bevacizumab-, temsirolimus-, or cytokine-containing regimens), or B.) Advanced renal cell carcinoma and used as first-line therapy in combination with avelumab or pembrolizumab |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

INQOVI

Products Affected

- INQOVI

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of myelodysplastic syndromes (MDS), including previously treated and untreated, de novo and secondary MDS with the following French-American-British subtypes (refractory anemia, refractory anemia with ringed sideroblasts, refractory anemia with excess blasts, and chronic myelomonocytic leukemia [CMML]) and intermediate-1, intermediate-2, and high-risk International Prognostic Scoring System groups |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

INREBIC

Products Affected

- INREBIC

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of intermediate-2 or high-risk primary or secondary (post-polycythemia vera or post-essential thrombocythemia) myelofibrosis (MF). |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

INTRAROSA

Products Affected

- INTRAROSA

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Vaginal bleeding or dysfunctional uterine bleeding of an undetermined origin, or B.) Known or suspected estrogen-dependent neoplasia |
| Required Medical Information | Diagnosis of one of the following A.) moderate to severe dyspareunia due to menopause, or B.) atrophic vaginitis due to menopause |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | Initial: 3 months, Renewal: 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

IRESSA

Products Affected

- IRESSA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of metastatic non-small cell lung cancer (NSCLC) and must meet all of the following 1.) Tumor has epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 (L858R) substitution mutations as detected by an FDA-approved test or Clinical Laboratory Improvement Amendments-approved facility and 2.) Used as first-line treatment |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ISTURISA

Products Affected

- ISTURISA

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of Cushing's disease in patients for whom pituitary surgery is not an option or has not been curative |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an endocrinologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ITRACONAZOLE

Products Affected

- *itraconazole oral*

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Ventricular dysfunction (e.g., congestive heart failure (CHF) or history of CHF), B.) Concurrent therapy with a CYP3A4 substrate (e.g., methadone, lovastatin, simvastatin, etc.), C.) Concurrent use of CYP2D6 inhibitors (e.g., bupropion, fluoxetine, paroxetine, quinidine, terbinafine), D.) Renal or hepatic impairment and concomitant use of colchicine, fesoterodine, solifenacin, or telithromycin, E.) Pregnancy |
| Required Medical Information | Diagnosis of one of the following A.) Systemic fungal infection (e.g., aspergillosis, histoplasmosis, blastomycosis), or B.) Onychomycosis confirmed by one of the following: positive potassium hydroxide (KOH) preparation, fungal culture, or nail biopsy |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 6 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ITRACONAZOLE SOLN

Products Affected

- *itraconazole oral*

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Ventricular dysfunction (e.g., congestive heart failure (CHF) or history of CHF), B.) Concurrent therapy with a CYP3A4 substrate (e.g., methadone, lovastatin, simvastatin, etc.), C.) Concurrent use of CYP2D6 inhibitors (e.g., bupropion, fluoxetine, paroxetine, quinidine, terbinafine), D.) Renal or hepatic impairment and concomitant use of colchicine, fesoterodine, solifenacin, or telithromycin, E.) Pregnancy |
| Required Medical Information | Diagnosis of candidiasis (esophageal or oropharyngeal) that is refractory to treatment with fluconazole |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 6 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

IVERMECTIN

Products Affected

- *ivermectin oral*

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | Prevention or treatment of COVID-19 |
| Required Medical Information | Diagnosis of one of the following: A.) Strongyloidiasis of the intestinal tract or B.) Onchocerciasis |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 1 month |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

JAKAFI

Products Affected

- JAKAFI

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Intermediate or high-risk myelofibrosis, including primary myelofibrosis, post-polycythemia vera myelofibrosis and post-essential thrombocythemia myelofibrosis, B.) Polycythemia vera AND patient has had an inadequate response to or is intolerant of hydroxyurea, C.) Acute graft versus host disease AND disease is refractory to steroid therapy, or D.) Chronic graft-versus-host disease (cGVHD) after failure of corticosteroid therapy (alone or in combination with a calcineurin inhibitor) and up to one additional line of systemic therapy |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 6 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

JUXTAPID

Products Affected

- JUXTAPID ORAL CAPSULE 10 MG, 20 MG, 30 MG, 5 MG

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Moderate to severe liver impairment or active liver disease including unexplained persistent abnormal liver function tests, B.) Pregnancy, or C.) Concomitant use with strong or moderate CYP3A4 inhibitors |
| Required Medical Information | Diagnosis of HoFH as confirmed by one of the following A.) Genetic confirmation of 2 mutations in the LDL receptor, ApoB, PCSK9, or LDL receptor adaptor protein 1 (LDLRAP1 or ARH), or B.) Both of the following 1.) Either untreated LDL-C greater than 500 mg/dL or treated LDL-C greater than 300 mg/dL, and 2.) Either xanthoma before 10 years of age or evidence of heterozygous FH in both parents |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | Initial: 6 months, Renewal: 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

KALYDECO

Products Affected

- KALYDECO

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of cystic fibrosis (CF) and the patient has 1 mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to ivacaftor potentiation based on clinical and/or in vitro assay data |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

KESIMPTA

Products Affected

- KESIMPTA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | Active Hepatitis B infection |
| Required Medical Information | Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a neurologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

KISQALI

Products Affected

- KISQALI (200 MG DOSE)
- KISQALI (400 MG DOSE)
- KISQALI (600 MG DOSE)

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of advanced or metastatic hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer and one of the following A.) The patient is a pre-or perimenopausal woman or man and the requested drug will be used in combination with an aromatase inhibitor as initial endocrine-based therapy, B.) The patient is a postmenopausal woman, the requested drug will be used in combination with an aromatase inhibitor as initial endocrine-based therapy, and the patient has experienced disease progression, an intolerable adverse event, or contraindication to Ibrance (palbociclib) or Verzenio (abemaciclib), C.) The patient is a pre-or perimenopausal woman or man and the requested drug is being used with fulvestrant as initial endocrine-based therapy, or D.) The patient is a postmenopausal woman, the requested drug is being used following disease progression on endocrine therapy, and the patient has experienced disease progression, an intolerable adverse event, or contraindication to Ibrance (palbociclib) or Verzenio (abemaciclib) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

KISQALI FEMARA

Products Affected

- KISQALI FEMARA (400 MG DOSE)
- KISQALI FEMARA (600 MG DOSE)
- KISQALI FEMARA(200 MG DOSE)

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of advanced or metastatic hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer and one of the following A.) The patient is pre-or perimenopausal woman or male and the requested drug will be used as initial endocrine-based therapy, B.) The patient is postmenopausal, the requested drug will be used as initial endocrine-based therapy, and the patient has experienced disease progression, an intolerable adverse event, or contraindication to Ibrance (palbociclib) or Verzenio (abemaciclib) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

KORLYM

Products Affected

- KORLYM

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) pregnancy, B.) coadministration with simvastatin, lovastatin, or CYP3A substrates with narrow therapeutic ranges, C.) concomitant treatment with systemic corticosteroids for serious medical conditions or illnesses, D.) history of unexplained vaginal bleeding, E.) endometrial hyperplasia with atypia or endometrial carcinoma |
| Required Medical Information | Diagnosis of endogenous Cushing syndrome in patients with type 2 diabetes mellitus or glucose intolerance and must meet all of the following 1.) Used to control hyperglycemia secondary to hypercortisolism, and 2.) Patient has failed or is not a candidate for surgery |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an endocrinologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

KOSELUGO

Products Affected

- KOSELUGO

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of neurofibromatosis type 1 (NF1) in a patient who has symptomatic, inoperable plexiform neurofibromas (PN) |
| Age Restrictions | 2 years of age to 17 years of age |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

KYNMOBI

Products Affected

- KYNMOBI

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Concurrent use with 5-HT(3) receptor antagonists (e.g. ondansetron, granisetron, dolasetron, palonosetron, alosetron etc.) |
| Required Medical Information | Diagnosis of Parkinson's disease (PD) and patient is experiencing acute intermittent hypomobility (defined as off episodes characterized by muscle stiffness, slow movements, or difficulty starting movements) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

LAPATINIB

Products Affected

- *lapatinib ditosylate*

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of advanced or metastatic breast cancer with tumors that overexpress human epidermal growth factor receptor 2 (HER2) AND meets one of the following A.) Used in combination with capecitabine in a patient who has received prior therapy including an anthracycline, a taxane, and trastuzumab, OR B.) Used in combination with letrozole in a postmenopausal female for whom hormonal therapy is indicated |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

LENALIDOMIDE

Products Affected

- *lenalidomide*

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | Pregnancy |
| Required Medical Information | Diagnosis of one of the following A.) Multiple myeloma and medication will be used in combination with dexamethasone, B.) Autologous hematopoietic stem-cell transplantation (HSCT) in multiple myeloma patients, C.) Transfusion-dependent anemia due to low- or intermediate-1-risk myelodysplastic syndrome (MDS) associated with a deletion 5q cytogenetic abnormality or without additional cytogenetic abnormalities, D.) Mantle cell lymphoma whose disease has relapsed or progressed after two prior therapies, one of which included bortezomib, E.) Follicular lymphoma and used in combination with rituximab, or F.) Marginal zone lymphoma and used in combination with rituximab |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

LENVIMA

Products Affected

- LENVIMA (10 MG DAILY DOSE)
- LENVIMA (12 MG DAILY DOSE)
- LENVIMA (14 MG DAILY DOSE)
- LENVIMA (18 MG DAILY DOSE)
- LENVIMA (20 MG DAILY DOSE)
- LENVIMA (24 MG DAILY DOSE)
- LENVIMA (4 MG DAILY DOSE)
- LENVIMA (8 MG DAILY DOSE)

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Locally recurrent or metastatic, progressive, radioactive iodine-refractory differentiated thyroid cancer, B.) Advanced renal cell carcinoma, in combination with everolimus, following one prior anti-angiogenic therapy, C.) Unresectable hepatocellular carcinoma, first-line therapy, D.) Advanced endometrial carcinoma that is not microsatellite instability-high or mismatch repair deficient, in combination with pembrolizumab, when disease has progressed following prior systemic therapy and patient is not a candidate for curative surgery or radiation, or E.) Advanced renal cell carcinoma, in combination with pembrolizumab and used as first-line therapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

LEUKINE

Products Affected

- LEUKINE INJECTION SOLUTION
RECONSTITUTED

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Patient has undergone allogeneic or autologous bone marrow transplant (BMT) and engraftment is delayed or failed, B.) Patient is undergoing autologous peripheral-blood progenitor cell transplant to mobilize progenitor cells for collection by leukapheresis, C.) Medication will be used for myeloid reconstitution after an autologous or allogeneic BMT, D.) Patient has acute myeloid leukemia and administration will be after completion of induction chemotherapy, E.) Hematopoietic subsyndrome of acute radiation syndrome (H-ARS) or F.) Autologous peripheral blood stem cell transplant, Following myeloablative chemotherapy. |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

LEUPROLIDE

Products Affected

- ELIGARD
- *leuprolide acetate injection*
- LUPRON DEPOT (1-MONTH)
- LUPRON DEPOT (3-MONTH)
- LUPRON DEPOT (4-MONTH)
- LUPRON DEPOT (6-MONTH)

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Pregnancy |
| Required Medical Information | Diagnosis of one of the following A.) Advanced or metastatic prostate cancer and patient has failed or is intolerant to Eligard (7.5 mg 1-month, 22.5 mg 3-month, 30 mg 4-month, & 45 mg 6-month depots only), B.) Endometriosis (3.75 mg 1-month & 11.25 mg 3-month depots only), C.) Anemia due to uterine leiomyomata (Fibroids) (3.75 mg 1-month & 11.25 mg 3-month depots only) and patient is preoperative, or D.) Central precocious puberty (idiopathic or neurogenic) in children |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | B vs D determination required per CMS guidance |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

LIDOCAINE PATCH

Products Affected

- *lidocaine external patch 5 %*

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Pain associated with diabetic neuropathy, B.) Pain associated with cancer-related neuropathy, C.) Post-herpetic neuralgia, D.) Chronic back pain, or E.) Osteoarthritis of the knee or hip |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

LINEZOLID

Products Affected

- *linezolid intravenous solution 600 mg/300ml*
- *linezolid oral tablet*

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Concomitant use of an MAOI, or B.) Use within 14 days of discontinuing an MAOI |
| Required Medical Information | Diagnosis of one of the following A.) Community acquired pneumonia, B.) Hospital-acquired pneumonia, C.) Vancomycin-resistant Enterococcus faecium infection, D.) Complicated skin and skin structure infections, or E.) Uncomplicated skin and skin structure infections |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 1 month |
| Other Criteria | B vs D determination required per CMS guidance |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

LIVMARLI

Products Affected

- LIVMARLI

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of cholestatic pruritus in patients with Alagille syndrome (ALGS) |
| Age Restrictions | 1 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

LIVTENCITY

Products Affected

- LIVTENCITY

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of post-transplant cytomegalovirus (CMV) infection/disease that is refractory to treatment (with or without genotypic resistance) with ganciclovir, valganciclovir, cidofovir or foscarnet |
| Age Restrictions | 12 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

LONSURF

Products Affected

- LONSURF

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Metastatic colorectal cancer, previously treated with fluoropyrimidine, oxaliplatin, and irinotecan-based regimens, an anti-VEGF therapy, and if RAS wild-type, an anti-EGFR therapy, or B.) Metastatic gastric or gastroesophageal junction adenocarcinoma previously treated with at least 2 prior lines of chemotherapy that included a fluoropyrimidine, a platinum, either a taxane or irinotecan and if appropriate, HER2/neu-targeted therapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

LORBRENA

Products Affected

- LORBRENA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | Concomitant use with strong CYP3A4 inducers |
| Required Medical Information | Diagnosis of anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC) as detected by an FDA-approved test |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

LUMAKRAS

Products Affected

- LUMAKRAS

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of KRAS G12C-mutated locally advanced or metastatic non-small cell lung cancer (NSCLC) as determined by an FDA-approved test and patient has received at least one prior systemic therapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

LUPKYNIS

Products Affected

- LUPKYNIS

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Concomitant use of strong CYP3A4 inhibitors (e.g., ketoconazole, itraconazole, clarithromycin) |
| Required Medical Information | Initial: Diagnosis of systemic lupus erythematosus (SLE) with active lupus nephritis (LN) Classes III, IV, V (alone or in combination), and all of the following: 1.) Baseline renal function of 45 mL/min/1.73 m ² or greater, 2.) Will be used in combination with a background immunosuppressive therapy regimen (e.g. mycophenolate, oral steroids, etc). Renewal: Improvement in urine protein to creatinine ratio (UPCR) (i.e. less than or equal to 0.5 mg/mg) AND estimated glomerular filtration rate (eGFR) of 60 mL/min/1.73 m ² or greater, or no confirmed decrease from baseline in eGFR of greater than 20% |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a rheumatologist or nephrologist |
| Coverage Duration | Initial: 12 months, Renewal: 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

LYNPARZA

Products Affected

- LYNPARZA ORAL TABLET

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) HER2-negative, deleterious or suspected deleterious germline BRCA mutated high-risk early or metastatic breast cancer AND patient has been previously treated with chemotherapy in neoadjuvant, adjuvant, or metastatic setting, B.) Advanced ovarian cancer with known or suspected BRCA mutation as detected by an FDA-approved test AND patient has trial and failure, contraindication, or intolerance to 3 or more prior lines of chemotherapy, C.) Recurrent epithelial ovarian cancer, recurrent fallopian tube cancer, or recurrent primary peritoneal cancer AND used for maintenance treatment in patients who are in complete or partial response to platinum-based chemotherapy (e.g. cisplatin, carboplatin), D.) Deleterious or suspected deleterious germline or somatic BRCA-mutated (gBRCAm or sBRCAm) epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients with complete or partial response to first-line platinum-based chemotherapy, E.) Deleterious or suspected deleterious germline BRCA-mutated metastatic pancreatic adenocarcinoma and disease has not progressed on at least 16 weeks of a first-line platinum-based chemotherapy regimen, F.) Advanced epithelial ovarian, fallopian tube, or primary peritoneal cancer in patients who are in complete or partial response to first-line platinum-based chemotherapy and whose cancer is associated with homologous recombination deficiency positive status defined by either a deleterious or suspected deleterious BRCA-mutation and/or genomic instability AND are using in combination with bevacizumab for maintenance treatment, or G.) Deleterious or suspected deleterious germline or somatic homologous recombination repair gene mutated metastatic castration-resistant prostate cancer in patients who have progressed following prior treatment with enzalutamide or abiraterone |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |

| PA Criteria | Criteria Details |
|---------------------|-------------------------------------|
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

MATULANE

Products Affected

- MATULANE

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | Inadequate marrow reserve |
| Required Medical Information | Diagnosis of Hodgkin's Disease, Stages III and IV and used in combination with other anticancer drugs |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

MAYZENT

Products Affected

- MAYZENT
- MAYZENT STARTER PACK

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) CYP2C9*3/*3 genotype, B.) In the last 6 months experienced myocardial infarction, unstable angina, stroke, TIA, decompensated heart failure requiring hospitalization, Class III-IV heart failure, or C.) Presence of Mobitz type II second-degree, third-degree AV block, or sick sinus syndrome, unless patient has a functioning pacemaker |
| Required Medical Information | Diagnosis of relapsing forms of multiple sclerosis, including clinically isolated syndrome, relapsing-remitting disease, or active secondary progressive disease and the following A.) Patients with relapsing forms of multiple sclerosis have history of/or contraindication to Avonex, Betaseron, Copaxone/Glatiramer, Gilenya/Fingolimod, or Dimethyl Fumarate |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a neurologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

MEKINIST

Products Affected

- MEKINIST

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Locally advanced or metastatic anaplastic thyroid cancer (ATC) with BRAF V600E mutation and used in combination with dabrafenib and no locoregional treatment options, B.) Malignant melanoma with lymph node involvement and following complete resection with BRAF V600E or V600K mutations and used in combination with dabrafenib, C.) Unresectable or metastatic malignant melanoma with BRAF V600E or V600K mutations and used in combination with dabrafenib or as monotherapy, D.) Metastatic non-small cell lung cancer, with BRAF V600E mutation, in combination with dabrafenib, or E.) Unresectable or metastatic solid tumors with BRAF V600E mutation, in combination with dabrafenib, and have progressed following prior treatment and have no satisfactory alternative treatment options |
| Age Restrictions | 6 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

MEKTOVI

Products Affected

- MEKTOVI

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of unresectable or metastatic malignant melanoma with documented BRAF V600E or V600K mutation as detected by an FDA approved test AND used in combination with encorafenib |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

MIGLUSTAT

Products Affected

- miglustat*

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of mild to moderate type 1 Gaucher disease and patient is not a candidate for enzyme replacement therapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

MS INTERFERONS

Products Affected

- AVONEX PEN INTRAMUSCULAR AUTO-INJECTOR KIT
- AVONEX PREFILLED INTRAMUSCULAR PREFILLED SYRINGE KIT
- BETASERON SUBCUTANEOUS KIT

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Relapsing forms of multiple sclerosis (e.g., clinically isolated syndrome, relapsing-remitting MS, active secondary progressive disease, or progressive-relapsing MS), or B.) Patient has experienced a first clinical episode and has MRI features consistent with multiple sclerosis |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a neurologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

NAMZARIC

Products Affected

- NAMZARIC

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Hypersensitivity to memantine, donepezil, or excipients, or B.) Hypersensitivity to piperidine derivatives |
| Required Medical Information | Diagnosis of moderate to severe dementia associated with Alzheimer's disease |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

NATPARA

Products Affected

- NATPARA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of hypoparathyroidism and used to control hypocalcemia |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

NERLYNX

Products Affected

- NERLYNX

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Early stage HER2-positive breast cancer and used following adjuvant trastuzumab therapy, or B.) Advanced or metastatic HER2-positive breast cancer, used in combination with capecitabine, AND patient has received 2 or more prior anti-HER2-based regimens in the metastatic setting |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

NINLARO

Products Affected

- NINLARO

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of multiple myeloma, used in combination with lenalidomide and dexamethasone, AND patient has history of at least 1 prior therapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

NITISINONE

Products Affected

- *nitisinone*
- ORFADIN ORAL CAPSULE 20 MG

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of hereditary tyrosinemia type 1 |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

NUBEQA

Products Affected

- NUBEQA

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Non-metastatic, castration-resistant prostate cancer, or B.) Metastatic hormone-sensitive prostate cancer in combination with docetaxel |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or urologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

NUCALA

Products Affected

- NUCALA

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Severe asthma with eosinophilic phenotype, B.) Eosinophilic granulomatosis with polyangiitis (EGPA), C.) Hypereosinophilic syndrome lasting at least 6 months without an identifiable non-hematologic secondary cause, or D.) Chronic rhinosinusitis with nasal polyps and used as an adjunct treatment |
| Age Restrictions | 6 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

NUEDEXTA

Products Affected

- NUEDEXTA

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) History of prolonged QT interval, congenital long QT syndrome or Torsades de pointes, B.) Heart failure, C.) Complete AV block without an implanted pacemaker or high risk of complete AV block, D.) Concomitant use with quinidine, quinine, mefloquine, or drugs that prolong QT interval and are metabolized by CYP2D6 (e.g., thioridazine, pimozide), E.) Concomitant use with MAOIs or within 14 days of MAOI therapy, F.) History of quinine-, mefloquine-, or quinidine-induced thrombocytopenia, bone marrow depression, or lupus-like syndrome |
| Required Medical Information | Diagnosis of pseudobulbar affect |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

NUPLAZID

Products Affected

- NUPLAZID ORAL CAPSULE
- NUPLAZID ORAL TABLET 10 MG

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of hallucinations and delusions associated with Parkinson disease psychosis |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

OCTREOTIDE

Products Affected

- *octreotide acetate injection solution 100 mcg/ml, 1000 mcg/ml, 200 mcg/ml, 50 mcg/ml, 500 mcg/ml*

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Acromegaly and patient has inadequate response to or is ineligible for surgery, radiation, or bromocriptine mesylate, or B.) Metastatic carcinoid syndrome, or C.) Vasoactive intestinal peptide-secreting tumors (VIPomas) with associated diarrhea |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | B vs D determination required per CMS guidance |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ODOMZO

Products Affected

- ODOMZO

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | Pregnancy |
| Required Medical Information | Diagnosis of locally advanced basal cell carcinoma of the skin and one of the following A.) Cancer has recurred following surgery or radiation therapy, B.) Patient is not a candidate for surgery or radiation therapy. |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

OFEV

Products Affected

- OFEV

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Idiopathic pulmonary fibrosis (IPF), B.) Systemic sclerosis-associated interstitial lung disease (ILD), or C.) Chronic fibrosing interstitial lung disease with a progressive phenotype |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a pulmonologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ONUREG

Products Affected

- ONUREG

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of acute myeloid leukemia (AML) used in maintenance treatment for adult patients who achieved first complete remission (CR) or complete remission with incomplete blood count recovery (CRi) following intensive induction chemotherapy and are not able to complete intensive curative therapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

OPSUMIT

Products Affected

- OPSUMIT

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | Pregnancy |
| Required Medical Information | Diagnosis of pulmonary arterial hypertension (WHO Group I), confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e. g., patient is frail, elderly, etc.) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a cardiologist or pulmonologist |
| Coverage Duration | Initial: 6 months, Renewal: 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ORGOVYX

Products Affected

- ORGOVYX

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of advanced prostate cancer |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or urologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ORKAMBI

Products Affected

- ORKAMBI

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of cystic fibrosis (CF) with documented homozygous F508del mutation confirmed by FDA-approved CF mutation test |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

OSPHERA

Products Affected

- OSPHERA

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Undiagnosed abnormal genital bleeding, B.) Known or suspected estrogen-dependent neoplasia, C.) Active deep vein thrombosis (DVT), pulmonary embolism (PE), or a history of these conditions, D.) Active arterial thromboembolic disease (e.g. stroke, myocardial infarction) or a history of these conditions, or E.) Pregnancy |
| Required Medical Information | Diagnosis of one of the following A.) Moderate to severe dyspareunia due to vulvar and vaginal atrophy associated with menopause, or B.) Moderate to severe vaginal dryness due to vulvar and vaginal atrophy associated with menopause |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

OXANDROLONE

Products Affected

- *oxandrolone oral*

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Known or suspected carcinoma of the prostate or breast in males, B.) Carcinoma of the breast in females with hypercalcemia, C.) Pregnancy, D.) Nephrosis or nephrotic phase of nephritis, E.) Hypercalcemia |
| Required Medical Information | Diagnosis of one of the following A.) Bone pain associated with osteoporosis, B.) Protein catabolism associated with chronic corticosteroid administration, or C.) Used as adjunctive therapy to promote weight gain after weight loss associated with one of the following 1.) Extensive surgery, 2.) Chronic infections, 3.) Severe trauma, or 4.) Failure to gain or maintain at least 90% of ideal body weight without definite pathophysiologic reasons |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 3 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

PANRETIN

Products Affected

- PANRETIN

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of AIDS-related Kaposi's sarcoma and both of the following 1.) Used to treat cutaneous lesions, and 2.) Systemic anti-Kaposi's Sarcoma therapy is not indicated (e.g., patient does not have more than 10 new KS lesions in the prior month, symptomatic lymphedema, symptomatic pulmonary KS, or symptomatic visceral involvement) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or HIV specialist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

PEGYLATED INTERFERON

Products Affected

- PEGASYS SUBCUTANEOUS SOLUTION 180 MCG/ML
- PEGASYS SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Autoimmune hepatitis, B.) Hepatic decompensation (Child-Pugh score greater than 6 (Class B and C) in cirrhotic patients before treatment, OR hepatic decompensation (Child-Pugh score greater than or equal to 6) in cirrhotic patients co-infected with hepatitis C and HIV before treatment, C.) Hypersensitivity reactions, including urticaria, bronchoconstriction, anaphylaxis, or Stevens-Johnson syndrome to alfa interferons or any component of the product, or D.) Pregnancy with concomitant ribavirin use |
| Required Medical Information | Diagnosis of one of the following A.) Chronic hepatitis B infection, or B.) Chronic hepatitis C and required criteria will be applied consistent with current AASLD-IDSA guidance with compensated liver disease |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with a gastroenterologist, hepatologist, or infectious disease specialist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

PEMAZYRE

Products Affected

- PEMAZYRE

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.)Previously treated, unresectable locally advanced or metastatic cholangiocarcinoma with confirmed fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement as detected by an FDA-approved test, or B.) Relapsed or refractory myeloid/lymphoid neoplasms with fibroblast growth factor receptor 1 rearrangement |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist, gastroenterologist, or hepatologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

PIQRAY

Products Affected

- PIQRAY (200 MG DAILY DOSE)
- PIQRAY (250 MG DAILY DOSE)
- PIQRAY (300 MG DAILY DOSE)

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of hormone receptor (HR) positive, HER2-negative, PIK3CA-mutated, advanced or metastatic breast cancer and must meet all of the following 1.) Used in combination with fulvestrant, 2.) Disease has progressed on or after an endocrine-based regimen, and 3.) Patient is a male or postmenopausal female |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

PIRFENIDONE

Products Affected

- *pirfenidone oral tablet*

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of idiopathic pulmonary fibrosis |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a pulmonologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

POMALYST

Products Affected

- POMALYST

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | Pregnancy |
| Required Medical Information | Diagnosis of one of the following A.) AIDS-related Kaposi sarcoma and patient has failure on highly active antiretroviral therapy (HAART), B.) Kaposi sarcoma in HIV-negative adults, or C.) Multiple myeloma and in combination with dexamethasone in adults who have received at least 2 prior therapies (including lenalidomide and a proteasome inhibitor) and have demonstrated disease progression on or within 60 days of completion of the last therapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

POSACONAZOLE

Products Affected

- *posaconazole*

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Concomitant treatment with sirolimus, B.) Concomitant use of CYP3A4 substrates that prolong QT interval (pimozide, quinidine), C.) Concomitant use of HMG-CoA Reductase inhibitors primarily metabolized through CYP3A4, or D.) Concomitant use of ergot alkaloids |
| Required Medical Information | Diagnosis of one of the following A.) Oropharyngeal candidiasis, B.) Patient is severely immunocompromised and requires prophylaxis of invasive aspergillosis or candidiasis due to high risk of infection, or C.) Invasive aspergillosis |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 weeks |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

POSACONAZOLE SUSPENSION

Products Affected

- NOXAFIL ORAL PACKET
- NOXAFIL ORAL SUSPENSION

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Concomitant treatment with sirolimus, B.) Concomitant use of CYP3A4 substrates that prolong QT interval (pimozide, quinidine), C.) Concomitant use of HMG-CoA Reductase inhibitors primarily metabolized through CYP3A4, or D.) Concomitant use of ergot alkaloids |
| Required Medical Information | Diagnosis of one of the following A.) Oropharyngeal candidiasis, B.) Patient is severely immunocompromised and requires prophylaxis of invasive aspergillosis or candidiasis due to high risk of infection, or C.) Invasive aspergillosis |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 weeks |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

PREVYMIS

Products Affected

- PREVYMIS ORAL

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Concomitant use with pimozide or ergot alkaloids (ergotamine, dihydroergotamine), B.) Concomitant use with pitavastatin or simvastatin when coadministered with cyclosporine |
| Required Medical Information | Must meet both of the following 1.) Patient is CMV-seropositive (R+) and 2.) Patient is receiving an allogeneic hematopoietic stem cell transplant (HSCT) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 6 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

PROMACTA

Products Affected

- PROMACTA

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Chronic idiopathic thrombocytopenic purpura (ITP), B.) Chronic hepatitis C infection associated thrombocytopenia, or C.) Severe aplastic anemia with insufficient response to immunosuppressive therapy or in combination with standard immunosuppressive therapy |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

QINLOCK

Products Affected

- QINLOCK

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of advanced gastrointestinal stromal tumor (GIST) and patient has received prior treatment with 3 or more kinase inhibitors, including imatinib |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

QUININE SULFATE

Products Affected

- *quinine sulfate oral*

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Prolongation of QT interval, B.) Glucose-6-phosphate dehydrogenase deficiency, C.) Myasthenia gravis, D.) Known hypersensitivity to mefloquine or quinidine, E.) Optic neuritis, F.) Diagnosis of Blackwater fever, G.) Use solely for treatment or prevention of nocturnal leg cramps |
| Required Medical Information | Diagnosis of one of the following A.) uncomplicated Plasmodium falciparum malaria, B.) uncomplicated Plasmodium vivax malaria, or C.) babesiosis |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 1 month |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

RAVICTI

Products Affected

- RAVICTI

| PA Criteria | Criteria Details |
|------------------------------|-------------------------------------|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of urea cycle disorders |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

REGRANEX

Products Affected

- REGRANEX

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | Known neoplasm at the site of application |
| Required Medical Information | Diagnosis of lower extremity diabetic neuropathic ulcers that extend into the subcutaneous tissue or beyond and have an adequate blood supply |
| Age Restrictions | 16 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

REPATHA

Products Affected

- REPATHA
- REPATHA SURECLICK
- REPATHA PUSHTRONEX SYSTEM

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) primary hyperlipidemia including heterozygous familial hypercholesterolemia (HeFH), B.) homozygous familial hypercholesterolemia, C.) established cardiovascular disease and myocardial infarction prophylaxis, stroke prophylaxis, or coronary revascularization prophylaxis is required, or D.) clinical atherosclerotic cardiovascular disease (CVD) as defined as one of the following 1.) acute coronary syndrome, 2.) history of myocardial infarction, 3.) stable/unstable angina, 4.) coronary or other arterial revascularization, 5.) stroke, 6.) transient ischemic stroke (TIA), or 7.) peripheral arterial disease presumed to be atherosclerotic region |
| Age Restrictions | 10 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

RETEVMO

Products Affected

- RETEVMO

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Advanced or metastatic RET-mutant medullary thyroid cancer (MTC) in patients who require systemic therapy, B.) Metastatic RET fusion-positive non-small cell lung cancer (NSCLC), C.) Advanced or metastatic RET fusion-positive thyroid cancer in patients who require systemic therapy and are refractory to radioactive iodine, if appropriate, or D.) Locally advanced or metastatic solid tumors with a RET gene fusion that have progressed on or following prior systemic treatment or who have no satisfactory alternative treatment options |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

REZUROCK

Products Affected

- REZUROCK

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of chronic graft-vs-host disease and patient has failed at least 2 prior lines of systemic therapy. |
| Age Restrictions | 12 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

RILUZOLE

Products Affected

- *riluzole*

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of amyotrophic lateral sclerosis (ALS) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

RINVOQ

Products Affected

- RINVOQ

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Moderate to severe rheumatoid arthritis, B.) Active psoriatic arthritis, C.) Moderate to severe atopic dermatitis and patient has trial/failure, contraindication, or intolerance to two of the following 1.) Topical corticosteroid and/or 2.) Topical calcineurin inhibitor, D.) Moderately to severely active ulcerative colitis who have had an inadequate response or intolerance to one or more tumor necrosis factor blockers, E.) Active ankylosing spondylitis who have had an inadequate response or intolerance to one or more tumor necrosis factor blockers, or F.) Active nonradiographic axial spondyloarthritis with objective signs of inflammation who have had an inadequate response or intolerance to tumor necrosis factor blocker therapy. |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | Screening for latent tuberculosis infection is required prior to initiation of treatment |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ROZLYTREK

Products Affected

- ROZLYTREK

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) ROS1-positive metastatic non-small cell lung cancer (NSCLC), or B.) Solid tumors that have a neurotrophic tyrosine receptor kinase (NTRK) gene fusion without a known acquired resistance mutation, are metastatic or where surgical resection is likely to result in severe morbidity, and have either progressed following treatment or have no satisfactory alternative therapy |
| Age Restrictions | 12 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

RUBRACA

Products Affected

- RUBRACA

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Epithelial ovarian, fallopian tube, or primary peritoneal cancer with deleterious BRCA mutation (germline and/or somatic) as detected by an FDA-approved test and patient has been treated with 2 or more prior lines of chemotherapy, B.) Recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer, used as maintenance treatment, and patient is in complete or partial response to platinum-based chemotherapy, or C.) Deleterious BRCA mutation (germline and/or somatic)-associated metastatic castration-resistant prostate cancer and patient has been treated with androgen receptor-directed therapy and a taxane-based chemotherapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

RYDAPT

Products Affected

- RYDAPT

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) treatment naive FLT3 mutation-positive acute myelogenous leukemia (AML) and must be used in combination with standard cytarabine and daunorubicin induction and cytarabine consolidation therapy, or B.) systemic mastocytosis or mast cell leukemia |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SAPROPTERIN

Products Affected

- sapropterin dihydrochloride oral packet*
- sapropterin dihydrochloride oral tablet*

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of hyperphenylalaninemia (HPA) caused by tetrahydrobiopterin (BH4)-responsive phenylketonuria (PKU) |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | Initial: 2 months, Renewal: 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SCSEMBLIX

Products Affected

- SCSEMBLIX

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP), previously treated with two or more tyrosine kinase inhibitors (TKIs), or B.) Philadelphia chromosome-positive chronic myeloid leukemia (Ph+ CML) in chronic phase (CP) with the T315I mutation |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SIGNIFOR

Products Affected

- SIGNIFOR

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of Cushing disease and patient has had inadequate response to or is not a candidate for surgery. For renewal: Documentation of a clinically meaningful reduction in 24-hour urinary free cortisol (UFC) levels or improvement in signs or symptoms of the disease |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | Initial: 6 months, Renewal: 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SILDENAFIL

Products Affected

- sildenafil citrate oral tablet 20 mg*

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Nitrate therapy, including intermittent use, B.) Concomitant use with riociguat or other guanylate cyclase stimulators, C.) Concomitant use with HIV protease inhibitors or elvitegravir/cobicistat/tenofovir/emtricitabine |
| Required Medical Information | Diagnosis of pulmonary arterial hypertension (WHO Group I), confirmed by right heart catheterization or Doppler echocardiogram if patient is unable to undergo a right heart catheterization (e.g., patient is frail, elderly, etc.) |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with a cardiologist or pulmonologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SIRTURO

Products Affected

- SIRTURO

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Must meet all of the following 1.) Diagnosis of pulmonary multidrug resistant tuberculosis (MDR-TB) and 2.) Used in combination with at least 3 other antibiotics for the treatment of pulmonary multi-drug resistant tuberculosis |
| Age Restrictions | 5 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an infectious disease specialist |
| Coverage Duration | 24 weeks |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SKYRIZI

Products Affected

- SKYRIZI PEN
- SKYRIZI SUBCUTANEOUS

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Moderate to severe plaque psoriasis and patient is a candidate for systemic therapy or phototherapy, B.) Active psoriatic arthritis, or C.) Moderately to severely active Crohn's disease in adults who have had an inadequate response to conventional therapy |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | Screening for latent tuberculosis infection is required prior to initiation of treatment |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SOLTAMOX

Products Affected

- SOLTAMOX

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Concomitant coumarin-type anticoagulant therapy, B.) history of thromboembolic disease such as DVT or PE |
| Required Medical Information | Diagnosis of breast cancer and documentation of inability to swallow tablet formulation |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SOMAVERT

Products Affected

- SOMAVERT

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of acromegaly and patient has had an inadequate response to or is ineligible for surgery or radiation therapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an endocrinologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SORAFENIB

Products Affected

- sorafenib tosylate*

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Squamous cell lung cancer being treated with carboplatin and paclitaxel |
| Required Medical Information | Diagnosis of one of the following A.) Advanced renal cell carcinoma, B.) Locally recurrent or metastatic, progressive, differentiated thyroid carcinoma that is refractory to radioactive iodine treatment, or C.) Unresectable hepatocellular carcinoma |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SPRYCEL

Products Affected

- SPRYCEL

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Newly diagnosed Philadelphia chromosome-positive (Ph+) chronic myeloid leukemia (CML) in chronic phase, B.) Chronic, accelerated, or myeloid or lymphoid blast phase Ph+ CML with resistance or intolerance to prior therapy, C.) Ph+ acute lymphoblastic leukemia (ALL) with resistance or intolerance to prior therapy, or D.) Newly diagnosed Ph+ ALL in combination with chemotherapy |
| Age Restrictions | 1 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

STELARA

Products Affected

- STELARA SUBCUTANEOUS SOLUTION 45 MG/0.5ML
- STELARA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Moderate to severely active Crohn disease, B.) Moderate to severe plaque psoriasis, C.) Active psoriatic arthritis, or D.) Moderate to severe active ulcerative colitis |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | Screening for latent tuberculosis infection is required prior to initiation of treatment |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

STIVARGA

Products Affected

- STIVARGA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Metastatic colorectal cancer in patients previously treated with fluoropyrimidine, oxaliplatin, and irinotecan containing chemotherapy, anti-VEGF therapy, and if RAS wild type, anti-EGFR therapy, B.) Liver carcinoma in patients previously treated with sorafenib, or C.) Locally advanced, unresectable or metastatic gastrointestinal stromal tumor (GIST) after treatment with imatinib and sunitinib |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SUNITINIB

Products Affected

- sunitinib malate*

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Gastrointestinal stromal tumor after disease progression on or intolerance to imatinib, B.) Pancreatic neuroendocrine tumors in a patient with unresectable locally advanced or metastatic disease, C.) Advanced renal cell carcinoma, or D.) Renal cell carcinoma and used as adjuvant therapy following nephrectomy in patients who are at high risk for recurrence |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SUNOSI

Products Affected

- SUNOSI

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Concomitant use of an MAOI, or B.) Use within 14 days of discontinuing an MAOI |
| Required Medical Information | Diagnosis of one of the following A.) narcolepsy with excessive daytime drowsiness and has trial of/or contraindication to modafinil or armodafinil, or B.) obstructive sleep apnea (OSA) with excessive daytime drowsiness and has trial of/or contraindication to modafinil or armodafinil |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SYMDEKO

Products Affected

- SYMDEKO

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of cystic fibrosis (CF) and must meet one of the following 1.) Patient is homozygous for the F508del mutation, or 2.) Patient has at least one mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene that is responsive to tezacaftor/ivacaftor verified by an FDA-cleared CF mutation test |
| Age Restrictions | 6 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation |
| Coverage Duration | Initial: 6 months, Renewal: 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SYMLIN

Products Affected

- SYMLINPEN 120 SUBCUTANEOUS SOLUTION PEN-INJECTOR
- SYMLINPEN 60 SUBCUTANEOUS SOLUTION PEN-INJECTOR

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Confirmed diagnosis of gastroparesis, B.) Hypoglycemia unawareness |
| Required Medical Information | Diagnosis of type 1 or type 2 diabetes mellitus and patient uses mealtime insulin therapy and has failed to achieve desired glucose control |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SYNAREL

Products Affected

- SYNAREL

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) pregnancy, B.) breastfeeding, C.) undiagnosed abnormal vaginal bleeding |
| Required Medical Information | Diagnosis of one of the following A.) Central precocious puberty, or B.) Endometriosis |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

SYNRIBO

Products Affected

- SYNRIBO

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of chronic or accelerated phase chronic myeloid leukemia (CML) and patient has tried and failed or has a contraindication or intolerance to at least 2 tyrosine kinase inhibitors |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | B vs D determination required per CMS guidance |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TABLOID

Products Affected

- TABLOID

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of acute myeloid leukemia (induction and consolidation therapy only) |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TABRECTA

Products Affected

- TABRECTA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of metastatic non-small cell lung cancer (NSCLC) in patients whose tumors have a mutation that leads to mesenchymal-epithelial transition (MET) exon 14 skipping as detected by an FDA-approved test |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TAFINLAR

Products Affected

- TAFINLAR

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Locally advanced or metastatic anaplastic thyroid carcinoma with BRAF V600E mutation, in combination with trametinib and no satisfactory locoregional treatment options, B.) Metastatic non-small cell lung cancer with BRAF V600E mutation, in combination with trametinib OR in patients previously treated as monotherapy, C.) Unresectable or metastatic malignant melanoma with BRAF V600E or V600K mutation, or D.) Unresectable or metastatic solid tumors with BRAF V600E mutation, in combination with trametinib, and have progressed following prior treatment and have no satisfactory alternative treatment options |
| Age Restrictions | 6 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TAGRISSO

Products Affected

- TAGRISSO

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Metastatic non-small cell lung cancer (NSCLC) with EGFR exon 19 deletion or exon 21 L858R mutation and used as first line therapy, B.) Metastatic non-small cell lung cancer with T790M EGFR mutation (as confirmed by an FDA-approved test) AND whose disease has progressed on or after EGFR tyrosine kinase inhibitor therapy, or C.) Non-small cell lung cancer (NSCLC) with tumor epidermal growth factor receptor (EGFR) exon 19 deletions or exon 21 L858R mutations (as confirmed by an FDA-approved test) AND patient requires adjuvant therapy after tumor resection |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TAKHZYRO

Products Affected

- TAKHZYRO

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following and used as routine prophylaxis A.) Hereditary angioedema (HAE) with C1 inhibitor deficiency (Type 1) confirmed by laboratory testing, or B.) HAE with C1 inhibitor dysfunction (Type 2) confirmed by laboratory testing, or C.) HAE with normal C1 inhibitor (Type 3) confirmed by laboratory testing and one of the following 1.) Positive test for an F12, angiopoietin-1, or plasminogen gene mutation, or 2.) Family history of angioedema and the angioedema was refractory to a trial of an antihistamine for at least one month |
| Age Restrictions | 12 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a hematologist, immunologist, or allergist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TALZENNA

Products Affected

- TALZENNA

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of deleterious or suspected deleterious germline breast cancer susceptibility gene (BRCA)-mutated (gBRCAm), human epidermal growth factor receptor 2 (HER2)-negative locally advanced or metastatic breast cancer |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TASIGNA

Products Affected

- TASIGNA

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Long QT syndrome, B.) Uncorrected hypokalemia, C.) Uncorrected hypomagnesemia |
| Required Medical Information | Diagnosis of one of the following A.) Newly diagnosed chronic phase Philadelphia chromosome-positive chronic myelogenous leukemia (CML), B.) Chronic phase or accelerated phase Philadelphia chromosome-positive CML in a patient with resistance or intolerance to prior therapy that included imatinib, or C.) Chronic phase Philadelphia chromosome-positive CML in a patient with resistance or intolerance to prior tyrosine-kinase inhibitor therapy |
| Age Restrictions | 1 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TAVNEOS

Products Affected

- TAVNEOS

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of severe active anti-neutrophil cytoplasmic autoantibody (ANCA)-associated vasculitis (granulomatosis with polyangiitis [GPA] and microscopic polyangiitis [MPA]) and both of the following apply 1.) Used as adjunctive treatment, and 2.) Used in combination with standard therapy including glucocorticoids |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TAZAROTENE

Products Affected

- *tazarotene external cream*
- *tazarotene external gel*
- TAZORAC EXTERNAL CREAM 0.05 %

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | Pregnancy |
| Required Medical Information | Diagnosis of one of the following A.) Acne vulgaris and patient has trial with at least one generic topical acne product, or B.) Stable moderate to severe plaque psoriasis with 20% or less body surface area involvement and patient has trial with at least one other topical psoriasis product (e.g., medium to high potency corticosteroid and/or vitamin D analogs) |
| Age Restrictions | 12 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TAZVERIK

Products Affected

- TAZVERIK

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Metastatic or locally advanced epithelioid sarcoma in patients not eligible for complete resection, B.) Relapsed or refractory follicular lymphoma in patients whose tumors are positive for an EZH2 mutation as detected by an FDA-approved test and who have received at least 2 prior systemic therapies, or C.) Relapsed or refractory follicular lymphoma in patients who have no satisfactory alternative treatment options |
| Age Restrictions | 16 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TEGSEDI

Products Affected

- TEGSEDI

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Platelet count less than 100,000 per microliter, B.) Urinary protein to creatinine ratio (UPCR) of 1000 mg/g or higher |
| Required Medical Information | Diagnosis of Polyneuropathy of hereditary transthyretin-mediated amyloidosis |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TEPMETKO

Products Affected

- TEPMETKO

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of metastatic non-small cell lung cancer (NSCLC) with mesenchymal-epithelial transition (MET) exon 14 skipping alterations |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TERIPARATIDE

Products Affected

- *teriparatide (recombinant)*

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Osteoporosis in postmenopausal female patient with high risk for fracture and patient has contraindication or has tried/had inadequate response to a bisphosphonate or Tymlos, B.) Primary or hypogonadal osteoporosis in male patient with high risk for fracture and patient has contraindication or has tried/had inadequate response to a bisphosphonate, or C.) Osteoporosis due to associated sustained systemic glucocorticoid therapy in patient with high risk for fracture and patient has contraindication or has tried/had inadequate response to a bisphosphonate |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | Initial: 12 months, Renewal: 12 months (Maximum 24 month treatment per patient lifetime) |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TETRABENAZINE

Products Affected

- tetrabenazine*

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Actively suicidal, B.) Untreated or inadequately treated depression, C.) Impaired hepatic function, D.) Concomitant use of monoamine oxidase inhibitors, E.) Concomitant use of reserpine or within 20 days of discontinuing reserpine |
| Required Medical Information | Diagnosis of chorea associated with Huntington's disease |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

THALOMID

Products Affected

- THALOMID

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | Pregnancy |
| Required Medical Information | Diagnosis of one of the following A.) Multiple myeloma that is newly diagnosed, or B.) Erythema nodosum leprosum (ENL) |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or infectious disease specialist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TIBSOVO

Products Affected

- TIBSOVO

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Relapsed or refractory acute myeloid leukemia with a susceptible isocitrate dehydrogenase-1 mutation (as detected by an FDA-approved test), B.) Previously treated, locally advanced or metastatic cholangiocarcinoma with an isocitrate dehydrogenase-1 mutation (as detected by an FDA-approved test.), or C.) Acute myeloid leukemia (newly-diagnosed) with susceptible isocitrate dehydrogenase-1 mutation and meets one of the following: 1.) Patient is 75 years of age or older, or 2.) Patient has comorbidities that preclude intensive induction chemotherapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a gastroenterologist, hematologist, hepatologist, or oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TOBI

Products Affected

- TOBI PODHALER

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | Known sensitivity to any aminoglycoside |
| Required Medical Information | Diagnosis of cystic fibrosis (confirmed by appropriate diagnostic or genetic testing) and patient has suspected or confirmed <i>Pseudomonas aeruginosa</i> infection in the lungs |
| Age Restrictions | 6 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TOLVAPTAN

Products Affected

- *tolvaptan*

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Diagnosis of Autosomal Dominant Polycystic Kidney Disease (ADPKD), B.) Urgent need to raise serum sodium acutely, C.) Inability to sense or appropriately respond to thirst, D.) Hypovolemic hyponatremia, E.) Concomitant use of strong CYP 3A Inhibitors (e.g. clarithromycin, ketoconazole, ritonavir), or F.) Anuria |
| Required Medical Information | Diagnosis of clinically significant hypervolemic or euvolemic hyponatremia (serum sodium less than 125 mEq/L or less marks hyponatremia that is symptomatic and has resisted correction with fluid restriction), including in patients with heart failure and syndrome of inappropriate antidiuretic hormone (SIADH) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 1 month |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TOPICAL RETINOIDS

Products Affected

- *tretinoin external*

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of mild to moderate acne vulgaris |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TOREMIFENE

Products Affected

- *toremifene citrate*

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Acquired or congenital long QT syndrome, B.) Uncorrected hypokalemia, C.) Uncorrected hypomagnesemia |
| Required Medical Information | Diagnosis of metastatic breast cancer and patient must have previous inadequate response or intolerance to tamoxifen |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 6 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TRELSTAR

Products Affected

- TRELSTAR MIXJECT

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of advanced prostate cancer and used in palliative treatment |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | B vs D determination required per CMS guidance |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TRIENTINE

Products Affected

- trientine hcl*

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of Wilson's disease in patients that are intolerant to penicillamine |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TRIKAFTA

Products Affected

- TRIKAFTA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of cystic fibrosis (CF) and patient has at least 1 F508del mutation in the cystic fibrosis transmembrane conductance regulator (CFTR) gene verified by an FDA-cleared CF mutation test |
| Age Restrictions | 6 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a pulmonologist or prescribing practitioner is from a CF center accredited by the Cystic Fibrosis Foundation |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TRUSELTIQ

Products Affected

- TRUSELTIQ (100MG DAILY DOSE)
- TRUSELTIQ (125MG DAILY DOSE)
- TRUSELTIQ (50MG DAILY DOSE)
- TRUSELTIQ (75MG DAILY DOSE)

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of previously treated, unresectable locally advanced or metastatic cholangiocarcinoma with a fibroblast growth factor receptor 2 (FGFR2) fusion or other rearrangement as detected by an FDA-approved test |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a gastroenterologist, hepatologist, or oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TUKYSA

Products Affected

- TUKYSA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of advanced unresectable or metastatic HER2-positive breast cancer (including brain metastases) in patients who have received one or more prior anti-HER2-based regimens in the metastatic setting and drug is being used in combination with trastuzumab and capecitabine |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TURALIO

Products Affected

- TURALIO ORAL CAPSULE 200 MG

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of symptomatic tenosynovial giant cell tumor (TGCT) associated with severe morbidity or functional limitations and not amenable to improvement with surgery |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

TYMLOS

Products Affected

- TYMLOS

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of osteoporosis in men or postmenopausal women and one of the following A.) osteoporotic fracture or multiple risk factors for fracture, or B.) previous trial of/or contraindication to bisphosphonate |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | Initial: 12 months, Renewal: 12 months (Maximum 24 month treatment per patient lifetime) |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

UBRELVY

Products Affected

- UBRELVY

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Concomitant use of strong CYP3A4 inhibitors (e.g., ketoconazole, itraconazole, clarithromycin) |
| Required Medical Information | Diagnosis of migraine disorder with or without aura and patient has documented trial, inadequate response, or contraindication to at least 1 generic formulary triptan |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

VALCHLOR

Products Affected

- VALCHLOR

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of cutaneous T-cell lymphoma (stage IA and IB mycosis fungoides-type) and patient has received prior skin-directed therapy (e.g. Topical corticosteroids, phototherapy, or topical nitrogen mustard) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

VENCLEXTA

Products Affected

- VENCLEXTA
- VENCLEXTA STARTING PACK

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Concomitant use with strong CYP3A inhibitor during the initial and titration phase in patients with CLL or SLL |
| Required Medical Information | Diagnosis of one of the following A.) chronic lymphocytic leukemia (CLL) or small lymphocytic lymphoma (SLL), or B.) Newly-diagnosed acute myeloid leukemia (AML) and used in combination with azacitidine, decitabine or low-dose cytarabine in patients 75 years or older or who have comorbidities that preclude use of intensive induction chemotherapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

VERQUVO

Products Affected

- VERQUVO

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Concomitant use of other soluble guanylate cyclase (sGC) stimulators, or B.) Pregnancy |
| Required Medical Information | Diagnosis of chronic heart failure (HF), NYHA Class II to IV and all of the following 1.) Left ventricular ejection fraction less than 45%, 2.) Previous hospitalization for HF within 6 months or outpatient IV diuretic treatment for HF within 3 months |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a cardiologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

VERZENIO

Products Affected

- VERZENIO

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative, node-positive, early breast cancer and ALL of the following: 1.) Patient is at high risk of recurrence, 2.) Ki-67 score 20% or greater as determined by an FDA approved test, and 3.) Requested drug will be used in combination with endocrine therapy (tamoxifen or an aromatase inhibitor) for adjuvant treatment, OR B.) Advanced or metastatic, hormone receptor (HR)-positive, human epidermal growth factor receptor 2 (HER2)-negative breast cancer and one of the following 1.) Used in combination with fulvestrant in a patient with disease progression following endocrine therapy, 2.) Used as monotherapy in a patient with disease progression following endocrine therapy and prior chemotherapy in the metastatic setting, or 3.) For postmenopausal women, and men, used as initial endocrine-based treatment in combination with an aromatase inhibitor |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

VIGABATRIN

Products Affected

- *vigabatrin*

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Infantile spasms, or B.) Refractory complex partial seizures and the drug is being used as adjunctive therapy in patients who have responded inadequately to two alternative treatments |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with a neurologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

VIJOICE

Products Affected

- VIJOICE

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of severe manifestations of PIK3CA-Related Overgrowth Spectrum (PROS) in patients who require systemic therapy |
| Age Restrictions | 2 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

VITRAKVI

Products Affected

- VITRAKVI

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of metastatic or surgically unresectable neurotrophic receptor tyrosine kinase (NTRK) gene fusion positive solid tumors without a known acquired resistance mutation and used in patients with unsatisfactory alternative treatments or who have progressed following treatment |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

VIZIMPRO

Products Affected

- VIZIMPRO

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of metastatic non-small cell lung cancer with confirmed epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 L858R substitution mutations as detected by an FDA-approved test |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

VONJO

Products Affected

- VONJO

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of intermediate or high-risk primary or secondary myelofibrosis in adults AND a platelet count less than 50 X 10(9)/L |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

VORICONAZOLE

Products Affected

- voriconazole intravenous
- voriconazole oral

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Concomitant use of carbamazepine, CYP3A4 substrates (e.g., terfenadine, astemizole, cisapride, pimozone, or quinidine), B.) Concomitant use with high-dose ritonavir (400mg every 12 hours), C.) Concomitant use with ergot alkaloids, D.) Concomitant use with long-acting barbiturates, E.) Concomitant use with rifabutin or rifampin, F.) Concomitant use with sirolimus, or G.) Concomitant use with efavirenz at standard doses of 400mg/day or higher |
| Required Medical Information | Diagnosis of one of the following A.) Invasive aspergillosis, B.) Candidemia, C.) Esophageal Candidiasis, D.) Invasive candidiasis of the skin and abdomen, kidney, bladder wall, and wounds, or E.) Serious fungal infection due to <i>Scedosporium apiospermum</i> or <i>Fusarium</i> species |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with an infectious disease specialist |
| Coverage Duration | 6 months |
| Other Criteria | IV formulation: B vs D determination required per CMS guidance |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

VOTRIENT

Products Affected

- VOTRIENT

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Advanced renal cell carcinoma, or B.) Advanced soft tissue sarcoma and patient received at least one prior chemotherapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

VYNDAMAX

Products Affected

- VYNDAMAX

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of wild type or hereditary transthyretin related familial amyloid cardiomyopathy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with a cardiologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

WELIREG

Products Affected

- WELIREG

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | Pregnancy |
| Required Medical Information | Diagnosis of von Hippel-Lindau (VHL) disease and therapy is required for any of the following disease associated tumors that do not require immediate surgery A.) Renal cell carcinoma (RCC), B.) Central nervous system (CNS) hemangioblastoma, or C.) Pancreatic neuroendocrine tumor (pNET) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

XALKORI

Products Affected

- XALKORI

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Metastatic non-small cell lung cancer (NSCLC) that is anaplastic lymphoma kinase (ALK)-positive or ROS1-positive as detected by an FDA-approved test, B.) Relapsed or refractory systemic anaplastic large cell lymphoma that is anaplastic lymphoma kinase (ALK) positive as detected by an FDA-approved test, or C.) Unresectable, recurrent, or refractory inflammatory myofibroblastic tumors that are anaplastic lymphoma kinase (ALK)-positive |
| Age Restrictions | None |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

XGEVA

Products Affected

- XGEVA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | Hypocalcemia (calcium less than 8.0 mg/dL) |
| Required Medical Information | Diagnosis of one of the following A.) Bone metastases from a solid tumor and used for the prevention of skeletal related events, B.) Multiple myeloma and used for the prevention of skeletal related events, C.) Hypercalcemia of malignancy refractory to bisphosphonate therapy, or D.) Giant cell tumor of bone that is unresectable or where surgical resection is likely to result in severe morbidity |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | B vs D determination required per CMS guidance |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

XOLAIR

Products Affected

- XOLAIR

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Chronic idiopathic urticaria in patients who remain symptomatic despite H1 antihistamine therapy and patient will continue to receive concurrent H1 antihistamine therapy unless contraindicated or not tolerated, B.) Moderate to severe persistent asthma in patients with a positive skin test or in vitro reactivity to a perennial aeroallergen and symptoms are inadequately controlled with inhaled corticosteroids and an additional controller medication (i.e. long acting beta2-agonist, leukotriene modifier, or sustained-release theophylline) and patient has trial and failure, contraindication, or intolerance to Dupixent or Nucala, or C.) Nasal polyps in patients with inadequate response to nasal corticosteroids, requested drug will be used as adjunctive treatment, and patient has trial and failure, contraindication, or intolerance to Dupixent |
| Age Restrictions | 6 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | B vs D determination required per CMS guidance |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

XOSPATA

Products Affected

- XOSPATA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of relapsed or refractory acute myeloid leukemia (AML) with a FMS-like tyrosine kinase 3 (FLT3) mutation as detected by an FDA-approved test |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

XPOVIO

Products Affected

- XPOVIO (100 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 50 MG
- XPOVIO (40 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 40 MG
- XPOVIO (40 MG TWICE WEEKLY) ORAL TABLET THERAPY PACK 40 MG
- XPOVIO (60 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 60 MG
- XPOVIO (60 MG TWICE WEEKLY)
- XPOVIO (80 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 40 MG
- XPOVIO (80 MG TWICE WEEKLY)

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Relapsed or refractory multiple myeloma being used in combination with dexamethasone in a patient who has received at least 4 prior therapies and is refractory to at least 2 proteasome inhibitors, at least 2 immunomodulatory agents, and an anti-CD38 monoclonal antibody, B.) Multiple myeloma being used in combination with bortezomib and dexamethasone in a patient who has received at least 1 prior therapy, C.) Relapsed or refractory diffuse large B-cell lymphoma not otherwise specified, or D.) Relapsed or refractory DLBCL arising from follicular lymphoma and patient has received at least 2 lines of systemic therapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

XTANDI

Products Affected

- XTANDI

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Castration-resistant prostate cancer, or B.) Metastatic, castration-sensitive prostate cancer |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or urologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

XURIDEN

Products Affected

- XURIDEN

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of hereditary orotic aciduria |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

XYREM

Products Affected

- XYREM

| PA Criteria | Criteria Details |
|-------------------------------------|--|
| Exclusion Criteria | Any of the following A.) Concomitant treatment with sedative hypnotic agents, B.) Succinic semialdehyde dehydrogenase deficiency |
| Required Medical Information | Diagnosis of one of the following A.) Narcolepsy with excessive daytime drowsiness and has trial of/or contraindication to a central nervous system (CNS) stimulant drug (e.g., amphetamine, dextroamphetamine, methylphenidate) or a CNS wakefulness promoting drug (e.g., armodafinil, modafinil), or B.) Cataplexy and narcolepsy |
| Age Restrictions | 7 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

XYWAV

Products Affected

- XYWAV

| PA Criteria | Criteria Details |
|-------------------------------------|---|
| Exclusion Criteria | Any of the following A.) Concomitant treatment with sedative hypnotic agents, B.) Succinic semialdehyde dehydrogenase deficiency |
| Required Medical Information | Diagnosis of one of the following A.) Narcolepsy with excessive daytime drowsiness and has trial of/or contraindication to a central nervous system (CNS) stimulant drug (e.g., amphetamine, dextroamphetamine, methylphenidate) or a CNS wakefulness promoting drug (e.g., armodafinil, modafinil), or B.) Cataplexy and narcolepsy, or C.) Idiopathic hypersomnia |
| Age Restrictions | 7 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

YONSA

Products Affected

- YONSA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | Pregnancy |
| Required Medical Information | Diagnosis of metastatic, castration-resistant prostate cancer and use in combination with methylprednisolone |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or urologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ZARXIO

Products Affected

- ZARXIO

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Chemotherapy induced febrile neutropenia (prophylaxis), B.) Severe chronic neutropenia, C.) Patient is undergoing autologous peripheral-blood progenitor cell transplant to mobilize progenitor cells for collection by leukapheresis, or D.) Hematopoietic subsyndrome of acute radiation syndrome (H-ARS) |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ZEJULA

Products Affected

- ZEJULA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Advanced or recurrent epithelial ovarian, fallopian tube, or primary peritoneal cancer and used as maintenance therapy in a patient who is in a complete or partial response to platinum-based chemotherapy, or B.) Advanced ovarian, fallopian tube, or primary peritoneal cancer and patient has been treated with 3 or more prior chemotherapy regimens, and cancer is associated with homologous recombination deficiency positive status defined by either a deleterious or suspected deleterious BRCA mutation, or genomic instability, and disease has progressed more than 6 months after response to the last platinum-based chemotherapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ZELBORAF

Products Affected

- ZELBORAF

| PA Criteria | Criteria Details |
|------------------------------|---|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of one of the following A.) Unresectable or metastatic melanoma and patient has positive BRAF-V600E mutation documented by an FDA-approved test, or B.) Erdheim-Chester disease and patient has documented BRAF V600 mutation |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ZIEXTENZO

Products Affected

- ZIEXTENZO

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of a non-myeloid malignancy and drug is being used as prophylaxis for chemotherapy-induced neutropenia |
| Age Restrictions | None |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ZOLINZA

Products Affected

- ZOLINZA

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of primary cutaneous T-cell lymphoma (CTCL) in patients who have progressive, persistent or recurrent disease on or following two systemic therapies (e.g., bexarotene, romidepsin, etc) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | None |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ZYDELIG

Products Affected

- ZYDELIG

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | History of toxic epidermal necrosis with any drug |
| Required Medical Information | Diagnosis of Chronic lymphocytic leukemia, used in combination with rituximab and patient has relapsed on at least one prior therapy |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist or hematologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

ZYKADIA

Products Affected

- ZYKADIA ORAL TABLET

| PA Criteria | Criteria Details |
|------------------------------|--|
| Exclusion Criteria | None |
| Required Medical Information | Diagnosis of anaplastic lymphoma kinase (ALK)-positive metastatic non-small cell lung cancer (NSCLC) |
| Age Restrictions | 18 years of age and older |
| Prescriber Restrictions | Prescribed by or in consultation with an oncologist |
| Coverage Duration | 12 months |
| Other Criteria | None |
| Indications | All Medically-accepted Indications. |
| Off-Label Uses | N/A |
| Part B Prerequisite | No |

PART B VERSUS PART D

This drug may be covered under Medicare Part B or D depending upon the circumstances. Information may need to be submitted describing the use and setting of the drug to make the determination.

Products Affected

- | | |
|--|---|
| • ABELCET INTRAVENOUS SUSPENSION 5 MG/ML | • cefoxitin sodium intravenous solution reconstituted 1 gm, 10 gm, 2 gm |
| • acetylcysteine inhalation solution 10 %, 20 % | • cefuroxime sodium injection solution reconstituted 750 mg |
| • acyclovir sodium intravenous solution 50 mg/ml | • cefuroxime sodium intravenous solution reconstituted 1.5 gm |
| • albuterol sulfate inhalation nebulization solution (2.5 mg/3ml) 0.083%, 0.63 mg/3ml, 1.25 mg/3ml, 2.5 mg/0.5ml | • chlorpromazine hcl oral tablet 10 mg, 25 mg |
| • AMBISOME INTRAVENOUS SUSPENSION RECONSTITUTED 50 MG | • cinacalcet hcl oral tablet 30 mg, 60 mg, 90 mg |
| • amikacin sulfate injection solution 500 mg/2ml | • ciprofloxacin in d5w intravenous solution 200 mg/100ml |
| • amphotericin b intravenous solution reconstituted 50 mg | • clindamycin phosphate injection solution 300 mg/2ml, 600 mg/4ml, 900 mg/6ml |
| • ampicillin sodium injection solution reconstituted 1 gm, 125 mg | • CLINIMIX E/DEXTROSE (2.75/5) INTRAVENOUS SOLUTION 2.75 % |
| • ampicillin sodium intravenous solution reconstituted 10 gm | • CLINIMIX E/DEXTROSE (4.25/10) INTRAVENOUS SOLUTION 4.25 % |
| • aprepitant oral capsule 125 mg, 40 mg, 80 & 125 mg, 80 mg | • CLINIMIX E/DEXTROSE (4.25/5) INTRAVENOUS SOLUTION 4.25 % |
| • AZASAN ORAL TABLET 100 MG, 75 MG | • CLINIMIX E/DEXTROSE (5/15) INTRAVENOUS SOLUTION 5 % |
| • azathioprine oral tablet 100 mg, 50 mg, 75 mg | • CLINIMIX E/DEXTROSE (5/20) INTRAVENOUS SOLUTION 5 % |
| • azithromycin intravenous solution reconstituted 500 mg | • CLINIMIX/DEXTROSE (4.25/10) INTRAVENOUS SOLUTION 4.25 % |
| • aztreonam injection solution reconstituted 2 gm | • CLINIMIX/DEXTROSE (4.25/5) INTRAVENOUS SOLUTION 4.25 % |
| • budesonide inhalation suspension 0.25 mg/2ml, 0.5 mg/2ml, 1 mg/2ml | • CLINIMIX/DEXTROSE (5/15) INTRAVENOUS SOLUTION 5 % |
| • calcitonin (salmon) nasal solution 200 unit/act | • CLINIMIX/DEXTROSE (5/20) INTRAVENOUS SOLUTION 5 % |
| • calcitriol oral capsule 0.25 mcg, 0.5 mcg | • colistimethate sodium (cba) injection solution reconstituted 150 mg |
| • calcitriol oral solution 1 mcg/ml | • cromolyn sodium inhalation nebulization solution 20 mg/2ml |
| • cefotetan disodium injection solution reconstituted 1 gm, 2 gm | • cyclophosphamide oral capsule 25 mg, 50 mg |

- cyclophosphamide oral tablet 25 mg, 50 mg
- cyclosporine modified oral capsule 100 mg, 25 mg, 50 mg
- cyclosporine modified oral solution 100 mg/ml
- cyclosporine oral capsule 100 mg, 25 mg
- dextrose intravenous solution 10 %, 5 %
- dextrose-nacl intravenous solution 10-0.2 %, 10-0.45 %, 2.5-0.45 %
- diphtheria-tetanus toxoids dt intramuscular suspension 25-5 lfu/0.5ml
- DOXY 100 INTRAVENOUS SOLUTION RECONSTITUTED 100 MG
- ENGERIX-B INJECTION SUSPENSION 20 MCG/ML
- ENGERIX-B INJECTION SUSPENSION PREFILLED SYRINGE 10 MCG/0.5ML, 20 MCG/ML
- ENVARSUS XR ORAL TABLET EXTENDED RELEASE 24 HOUR 0.75 MG, 1 MG, 4 MG
- ERAXIS INTRAVENOUS SOLUTION RECONSTITUTED 100 MG, 50 MG
- ERYTHROCIN LACTOBIONATE INTRAVENOUS SOLUTION RECONSTITUTED 500 MG
- everolimus oral tablet 0.25 mg, 0.5 mg, 0.75 mg, 1 mg
- fluconazole in sodium chloride intravenous solution 200-0.9 mg/100ml-%, 400-0.9 mg/200ml-%
- furosemide injection solution 10 mg/ml
- GENGRAF ORAL CAPSULE 100 MG, 25 MG
- GENGRAF ORAL SOLUTION 100 MG/ML
- granisetron hcl oral tablet 1 mg
- heparin sodium (porcine) injection solution 1000 unit/ml, 10000 unit/ml, 20000 unit/ml, 5000 unit/ml
- IMOVAX RABIES INTRAMUSCULAR SUSPENSION RECONSTITUTED 2.5 UNIT/ML
- INTRALIPID INTRAVENOUS EMULSION 20 %, 30 %
- ipratropium bromide inhalation solution 0.02 %
- ipratropium-albuterol inhalation solution 0.5-2.5 (3) mg/3ml
- ISOLYTE-P IN D5W INTRAVENOUS SOLUTION
- ISOLYTE-S PH 7.4 INTRAVENOUS SOLUTION
- kcl in dextrose-nacl intravenous solution 10-5-0.45 meq/l-%-%, 20-5-0.2 meq/l-%-%, 20-5-0.45 meq/l-%-%, 20-5-0.9 meq/l-%-%, 30-5-0.45 meq/l-%-%, 40-5-0.45 meq/l-%-%, 40-5-0.9 meq/l-%-%
- kcl-lactated ringers-d5w intravenous solution 20 meq/l
- methotrexate sodium (pf) injection solution 50 mg/2ml
- methotrexate sodium injection solution 50 mg/2ml
- methotrexate sodium oral tablet 2.5 mg
- methylprednisolone oral tablet 16 mg, 32 mg, 4 mg, 8 mg
- metronidazole intravenous solution 500 mg/100ml
- moxifloxacin hcl in nacl intravenous solution 400 mg/250ml
- mycophenolate mofetil oral capsule 250 mg
- mycophenolate mofetil oral suspension reconstituted 200 mg/ml
- mycophenolate mofetil oral tablet 500 mg
- mycophenolate sodium oral tablet delayed release 180 mg, 360 mg
- nafcillin sodium injection solution reconstituted 1 gm, 2 gm
- nafcillin sodium intravenous solution reconstituted 10 gm
- NUTRILIPID INTRAVENOUS EMULSION 20 %
- ondansetron hcl oral solution 4 mg/5ml
- ondansetron hcl oral tablet 4 mg, 8 mg
- ondansetron oral tablet dispersible 4 mg, 8 mg
- oxacillin sodium in dextrose intravenous solution 1 gm/50ml, 2 gm/50ml

March 2023

H9306_23_DRS_006_OE_C

- *oxacillin sodium injection solution reconstituted 1 gm, 2 gm*
- *oxacillin sodium intravenous solution reconstituted 10 gm*
- PANZYGA INTRAVENOUS SOLUTION 1 GM/10ML, 10 GM/100ML, 2.5 GM/25ML, 20 GM/200ML, 30 GM/300ML, 5 GM/50ML
- *paricalcitol oral capsule 1 mcg, 2 mcg, 4 mcg*
- *penicillin g potassium injection solution reconstituted 20000000 unit*
- *penicillin g sodium injection solution reconstituted 5000000 unit*
- *pentamidine isethionate inhalation solution reconstituted 300 mg*
- *pentamidine isethionate injection solution reconstituted 300 mg*
- *perphenazine oral tablet 4 mg, 8 mg*
- PLASMA-LYTE 148 INTRAVENOUS SOLUTION
- PLASMA-LYTE A INTRAVENOUS SOLUTION
- *potassium chloride in nacl intravenous solution 20-0.45 meq/l-%, 20-0.9 meq/l-%, 40-0.9 meq/l-%*
- *potassium chloride intravenous solution 2 meq/ml, 2 meq/ml (20 ml), 20 meq/100ml*
- *potassium cl in dextrose 5% intravenous solution 20 meq/l*
- *prednisolone oral solution 15 mg/5ml*
- *prednisolone sodium phosphate oral solution 10 mg/5ml, 20 mg/5ml, 25 mg/5ml, 6.7 (5 base) mg/5ml*
- *prednisolone sodium phosphate oral tablet dispersible 10 mg, 15 mg, 30 mg*
- PREDNISON INTENSOL ORAL CONCENTRATE 5 MG/ML
- *prednisone oral solution 5 mg/5ml*
- *prednisone oral tablet 1 mg, 10 mg, 2.5 mg, 20 mg, 5 mg, 50 mg*
- *prehevbrio intramuscular suspension 10 mcg/ml*
- PREMASOL INTRAVENOUS SOLUTION 10 %
- PRIVIGEN INTRAVENOUS SOLUTION 20 GM/200ML
- *prochlorperazine maleate oral tablet 10 mg, 5 mg*
- PROGRAF ORAL PACKET 0.2 MG, 1 MG
- PROSOL INTRAVENOUS SOLUTION 20 %
- PULMOZYME INHALATION SOLUTION 2.5 MG/2.5ML
- RABAVERT INTRAMUSCULAR SUSPENSION RECONSTITUTED
- RECOMBIVAX HB INJECTION SUSPENSION 10 MCG/ML, 40 MCG/ML, 5 MCG/0.5ML
- RECOMBIVAX HB INJECTION SUSPENSION PREFILLED SYRINGE 10 MCG/ML, 5 MCG/0.5ML
- *sirolimus oral solution 1 mg/ml*
- *sirolimus oral tablet 0.5 mg, 1 mg, 2 mg*
- *tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg*
- TDVAX INTRAMUSCULAR SUSPENSION 2-2 LF/0.5ML
- TEFLARO INTRAVENOUS SOLUTION RECONSTITUTED 400 MG, 600 MG
- TENIVAC INTRAMUSCULAR INJECTABLE 5-2 LFU, 5-2 LFU (INJECTION)
- *tigecycline intravenous solution reconstituted 50 mg*
- *tobramycin inhalation nebulization solution 300 mg/5ml*
- *tobramycin sulfate injection solution 10 mg/ml, 80 mg/2ml*
- TPN ELECTROLYTES INTRAVENOUS CONCENTRATE
- TRAVASOL INTRAVENOUS SOLUTION 10 %
- TREXALL ORAL TABLET 10 MG, 15 MG, 5 MG, 7.5 MG
- TROPHAMINE INTRAVENOUS SOLUTION 10 %
- VARUBI (180 MG DOSE) ORAL TABLET THERAPY PACK 2 X 90 MG

March 2023

H9306_23_DRS_006_OE_C

- XATMEP ORAL SOLUTION 2.5
MG/ML

Index

A

| | |
|---|-----|
| ABELCET INTRAVENOUS SUSPENSION 5 MG/ML..... | 245 |
| abiraterone acetate | 1 |
| acetylcysteine inhalation solution 10 %, 20 % | 245 |
| acitretin | 2 |
| ACTIMMUNE..... | 3 |
| acyclovir sodium intravenous solution 50 mg/ml | 245 |
| ADEMPAS | 4 |
| albuterol sulfate inhalation nebulization solution (2.5 mg/3ml) 0.083%, 0.63 mg/3ml, 1.25 mg/3ml, 2.5 mg/0.5ml .. | 245 |
| ALECENSA..... | 5 |
| ALUNBRIG..... | 7 |
| AMBISOME INTRAVENOUS SUSPENSION RECONSTITUTED 50 MG | 245 |
| ambrisentan | 8 |
| amikacin sulfate injection solution 500 mg/2ml | 245 |
| amphotericin b intravenous solution reconstituted 50 mg..... | 245 |
| ampicillin sodium injection solution reconstituted 1 gm, 125 mg..... | 245 |
| ampicillin sodium intravenous solution reconstituted 10 gm..... | 245 |
| aprepitant oral capsule 125 mg, 40 mg, 80 & 125 mg, 80 mg | 245 |
| ARCALYST | 9 |
| ARIKAYCE..... | 10 |
| armodafinil..... | 31 |
| AURYXIA | 11 |
| AUSTEDO..... | 12 |
| AVONEX PEN INTRAMUSCULAR AUTO-INJECTOR KIT..... | 128 |
| AVONEX PREFILLED INTRAMUSCULAR PREFILLED SYRINGE KIT..... | 128 |
| AYVAKIT | 13 |
| AZASAN ORAL TABLET 100 MG, 75 MG | 245 |

| | |
|--|-----|
| azathioprine oral tablet 100 mg, 50 mg, 75 mg | 245 |
| azithromycin intravenous solution reconstituted 500 mg..... | 245 |
| aztreonam injection solution reconstituted 2 gm | 245 |

B

| | |
|---|-----|
| BALVERSA | 14 |
| BENLYSTA SUBCUTANEOUS..... | 15 |
| BESREMI | 16 |
| BETASERON SUBCUTANEOUS KIT | 128 |
| bexarotene external | 17 |
| bexarotene oral..... | 18 |
| bosentan | 19 |
| BOSULIF | 20 |
| BRAFTOVI ORAL CAPSULE 75 MG ... | 21 |
| BRUKINSA | 22 |
| budesonide inhalation suspension 0.25 mg/2ml, 0.5 mg/2ml, 1 mg/2ml | 245 |
| BYLVAY | 23 |
| BYLVAY (PELLETS) ORAL CAPSULE SPRINKLE 200 MCG | 23 |

C

| | |
|---|-----|
| CABLIVI | 24 |
| CABOMETYX | 25 |
| calcitonin (salmon) nasal solution 200 unit/act..... | 245 |
| calcitriol oral capsule 0.25 mcg, 0.5 mcg | 245 |
| calcitriol oral solution 1 mcg/ml | 245 |
| CALQUENCE | 26 |
| CAMZYOS | 27 |
| CAPRELSA | 28 |
| carglumic acid oral tablet soluble | 29 |
| CAYSTON..... | 30 |
| cefotetan disodium injection solution reconstituted 1 gm, 2 gm..... | 245 |
| cefoxitin sodium intravenous solution reconstituted 1 gm, 10 gm, 2 gm..... | 245 |
| cefuroxime sodium injection solution reconstituted 750 mg..... | 245 |
| cefuroxime sodium intravenous solution reconstituted 1.5 gm..... | 245 |

| | |
|---|-----|
| chlorpromazine hcl oral tablet 10 mg, 25 mg | 245 |
| cinacalcet hcl oral tablet 30 mg, 60 mg, 90 mg | 245 |
| ciprofloxacin in d5w intravenous solution 200 mg/100ml | 245 |
| clindamycin phosphate injection solution 300 mg/2ml, 600 mg/4ml, 900 mg/6ml | 245 |
| CLINIMIX E/DEXTROSE (2.75/5) INTRAVENOUS SOLUTION 2.75 % | 245 |
| CLINIMIX E/DEXTROSE (4.25/10) INTRAVENOUS SOLUTION 4.25 % | 245 |
| CLINIMIX E/DEXTROSE (4.25/5) INTRAVENOUS SOLUTION 4.25 % | 245 |
| CLINIMIX E/DEXTROSE (5/15) INTRAVENOUS SOLUTION 5 %.... | 245 |
| CLINIMIX E/DEXTROSE (5/20) INTRAVENOUS SOLUTION 5 %.... | 245 |
| CLINIMIX/DEXTROSE (4.25/10) INTRAVENOUS SOLUTION 4.25 % | 245 |
| CLINIMIX/DEXTROSE (4.25/5) INTRAVENOUS SOLUTION 4.25 % | 245 |
| CLINIMIX/DEXTROSE (5/15) INTRAVENOUS SOLUTION 5 %.... | 245 |
| CLINIMIX/DEXTROSE (5/20) INTRAVENOUS SOLUTION 5 %.... | 245 |
| colistimethate sodium (cba) injection solution reconstituted 150 mg | 245 |
| COMETRIQ (100 MG DAILY DOSE) ORAL KIT 80 & 20 MG | 32 |
| COMETRIQ (140 MG DAILY DOSE) ORAL KIT 3 X 20 MG & 80 MG | 32 |
| COMETRIQ (60 MG DAILY DOSE)..... | 32 |
| COPAXONE SUBCUTANEOUS SOLUTION PREFILLED SYRINGE .. | 75 |
| COPIKTRA..... | 33 |
| CORLANOR ORAL TABLET | 34 |
| COSENTYX (300 MG DOSE)..... | 35 |
| COSENTYX SENSOREADY (300 MG). 35 | |
| COSENTYX SUBCUTANEOUS SOLUTION PREFILLED SYRINGE 75 MG/0.5ML | 35 |
| COTELLIC | 36 |
| cromolyn sodium inhalation nebulization solution 20 mg/2ml | 245 |

| | |
|---|-----|
| cyclophosphamide oral capsule 25 mg, 50 mg | 245 |
| cyclophosphamide oral tablet 25 mg, 50 mg | 246 |
| cyclosporine modified oral capsule 100 mg, 25 mg, 50 mg | 246 |
| cyclosporine modified oral solution 100 mg/ml | 246 |
| cyclosporine oral capsule 100 mg, 25 mg | 246 |
| CYSTADROPS..... | 38 |
| CYSTAGON..... | 37 |
| CYSTARAN | 38 |
| D | |
| dalfampridine er | 39 |
| DAURISMO | 40 |
| deferasirox granules | 41 |
| deferasirox oral tablet | 41 |
| deferasirox oral tablet soluble..... | 41 |
| deferiprone | 42 |
| dextrose intravenous solution 10 %, 5 % | 246 |
| dextrose-nacl intravenous solution 10-0.2 %, 10-0.45 %, 2.5-0.45 % | 246 |
| DIACOMIT..... | 43 |
| diclofenac sodium external gel 3 % | 44 |
| DIFICID..... | 45 |
| dimethyl fumarate oral..... | 46 |
| dimethyl fumarate starter pack..... | 46 |
| diphtheria-tetanus toxoids dt intramuscular suspension 25-5 lfu/0.5ml | 246 |
| DOJOLVI..... | 47 |
| DOXY 100 INTRAVENOUS SOLUTION RECONSTITUTED 100 MG | 246 |
| dronabinol | 48 |
| droxidopa | 49 |
| DUPIXENT..... | 50 |
| E | |
| ELIGARD | 112 |
| EMGALITY | 51 |
| ENBREL MINI..... | 52 |
| ENBREL SUBCUTANEOUS SOLUTION 25 MG/0.5ML | 52 |
| ENBREL SUBCUTANEOUS SOLUTION PREFILLED SYRINGE | 52 |
| ENBREL SURECLICK SUBCUTANEOUS SOLUTION AUTO-INJECTOR | 52 |
| ENDARI | 53 |

| | |
|--|-----|
| ENGERIX-B INJECTION SUSPENSION 20 MCG/ML | 246 |
| ENGERIX-B INJECTION SUSPENSION PREFILLED SYRINGE 10 MCG/0.5ML, 20 MCG/ML | 246 |
| ENSPRYNG | 54 |
| ENVARUSUS XR ORAL TABLET EXTENDED RELEASE 24 HOUR 0.75 MG, 1 MG, 4 MG | 246 |
| EPIDIOLEX..... | 55 |
| ERAXIS INTRAVENOUS SOLUTION RECONSTITUTED 100 MG, 50 MG | 246 |
| ERIVEDGE..... | 57 |
| ERLEADA..... | 58 |
| erlotinib hcl | 59 |
| ERYTHROCIN LACTOBIONATE INTRAVENOUS SOLUTION RECONSTITUTED 500 MG | 246 |
| ESBRIET ORAL CAPSULE..... | 60 |
| everolimus oral tablet 0.25 mg, 0.5 mg, 0.75 mg, 1 mg | 246 |
| everolimus oral tablet 10 mg, 2.5 mg, 5 mg, 7.5 mg | 61 |
| everolimus oral tablet soluble | 62 |
| EVRYSDI | 63 |
| EXKIVITY | 64 |
| F | |
| febuxostat..... | 65 |
| fentanyl | 67 |
| fentanyl citrate buccal lozenge on a handle | 66 |
| FERRIPROX ORAL SOLUTION..... | 42 |
| FERRIPROX TWICE-A-DAY..... | 42 |
| fingolimod hcl..... | 68 |
| FINTEPLA..... | 69 |
| FIRAZYR | 84 |
| fluconazole in sodium chloride intravenous solution 200-0.9 mg/100ml-%, 400-0.9 mg/200ml-% | 246 |
| FOTIVDA | 70 |
| furosemide injection solution 10 mg/ml . | 246 |
| G | |
| GALAFOLD | 71 |
| GATTEX..... | 72 |
| GAVRETO | 73 |

| | |
|---|--------|
| GENGRAF ORAL CAPSULE 100 MG, 25 MG | 246 |
| GENGRAF ORAL SOLUTION 100 MG/ML | 246 |
| GILOTRIF | 74 |
| GLEOSTINE ORAL CAPSULE 10 MG, 100 MG, 40 MG..... | 76 |
| granisetron hcl oral tablet 1 mg | 246 |
| H | |
| heparin sodium (porcine) injection solution 1000 unit/ml, 10000 unit/ml, 20000 unit/ml, 5000 unit/ml | 246 |
| HUMIRA PEDIATRIC CROHNS START SUBCUTANEOUS PREFILLED SYRINGE KIT 80 MG/0.8ML, 80 MG/0.8ML & 40MG/0.4ML | 80, 81 |
| HUMIRA PEN SUBCUTANEOUS PEN- INJECTOR KIT | 80, 81 |
| HUMIRA PEN-CD/UC/HS STARTER .. | 80, 81 |
| HUMIRA PEN-PEDIATRIC UC START | 80, 81 |
| HUMIRA PEN-PS/UV/ADOL HS START SUBCUTANEOUS PEN-INJECTOR KIT 40 MG/0.8ML | 80, 81 |
| HUMIRA PEN-PSOR/UEIT STARTER | 80, 81 |
| HUMIRA SUBCUTANEOUS PREFILLED SYRINGE KIT 10 MG/0.1ML, 20 MG/0.2ML, 40 MG/0.4ML, 40 MG/0.8ML | 80, 81 |
| HYFTOR..... | 82 |
| I | |
| IBRANCE | 83 |
| icatibant acetate..... | 84 |
| ICLUSIG..... | 85 |
| IDHIFA | 86 |
| imatinib mesylate | 87 |
| IMBRUVICA..... | 88 |
| IMOVAX RABIES INTRAMUSCULAR SUSPENSION RECONSTITUTED 2.5 UNIT/ML | 246 |
| INCRELEX..... | 89 |
| INLYTA..... | 90 |
| INQOVI | 91 |
| INREBIC..... | 92 |

| | |
|---|--------|
| INTRALIPID INTRAVENOUS | |
| EMULSION 20 %, 30 % | 246 |
| INTRAROSA..... | 93 |
| ipratropium bromide inhalation solution | |
| 0.02 % | 246 |
| ipratropium-albuterol inhalation solution | |
| 0.5-2.5 (3) mg/3ml | 246 |
| IRESSA..... | 94 |
| ISOLYTE-P IN D5W INTRAVENOUS | |
| SOLUTION..... | 246 |
| ISOLYTE-S PH 7.4 INTRAVENOUS | |
| SOLUTION..... | 246 |
| ISTURISA..... | 95 |
| itraconazole oral..... | 96, 97 |
| ivermectin oral | 98 |
| J | |
| JAKAFI..... | 99 |
| JUXTAPID ORAL CAPSULE 10 MG, 20 | |
| MG, 30 MG, 5 MG | 100 |
| K | |
| KALYDECO..... | 101 |
| kcl in dextrose-nacl intravenous solution 10- | |
| 5-0.45 meq/l-%-%, 20-5-0.2 meq/l-%-%, | |
| 20-5-0.45 meq/l-%-%, 20-5-0.9 meq/l-%- | |
| %, 30-5-0.45 meq/l-%-%, 40-5-0.45 | |
| meq/l-%-%, 40-5-0.9 meq/l-%-% | 246 |
| kcl-lactated ringers-d5w intravenous | |
| solution 20 meq/l..... | 246 |
| KESIMPTA..... | 102 |
| KISQALI (200 MG DOSE)..... | 103 |
| KISQALI (400 MG DOSE)..... | 103 |
| KISQALI (600 MG DOSE)..... | 103 |
| KISQALI FEMARA (400 MG DOSE) .. | 104 |
| KISQALI FEMARA (600 MG DOSE) .. | 104 |
| KISQALI FEMARA(200 MG DOSE) ... | 104 |
| KORLYM | 105 |
| KOSELUGO | 106 |
| KYNMOBI | 107 |
| L | |
| lapatinib ditosylate..... | 108 |
| lenalidomide..... | 109 |
| LENVIMA (10 MG DAILY DOSE) | 110 |
| LENVIMA (12 MG DAILY DOSE) | 110 |
| LENVIMA (14 MG DAILY DOSE) | 110 |
| LENVIMA (18 MG DAILY DOSE) | 110 |
| LENVIMA (20 MG DAILY DOSE) | 110 |

| | |
|---|----------|
| LENVIMA (24 MG DAILY DOSE) | 110 |
| LENVIMA (4 MG DAILY DOSE) | 110 |
| LENVIMA (8 MG DAILY DOSE) | 110 |
| LEUKINE INJECTION SOLUTION | |
| RECONSTITUTED..... | 111 |
| leuprolide acetate injection | 112 |
| lidocaine external patch 5 % | 113 |
| linezolid intravenous solution 600 mg/300ml | |
| | 114 |
| linezolid oral tablet | 114 |
| LIVMARLI..... | 115 |
| LIVTENCITY..... | 116 |
| LONSURF | 117 |
| LORBRENA..... | 118 |
| LUMAKRAS | 119 |
| LUPKYNIS..... | 120 |
| LUPRON DEPOT (1-MONTH) | 112 |
| LUPRON DEPOT (3-MONTH) | 112 |
| LUPRON DEPOT (4-MONTH) | 112 |
| LUPRON DEPOT (6-MONTH) | 112 |
| LYNPARZA ORAL TABLET | 121, 122 |
| M | |
| MATULANE | 123 |
| MAVYRET..... | 79 |
| MAYZENT | 124 |
| MAYZENT STARTER PACK..... | 124 |
| MEKINIST | 125 |
| MEKTOVI | 126 |
| methotrexate sodium (pf) injection solution | |
| 50 mg/2ml | 246 |
| methotrexate sodium injection solution 50 | |
| mg/2ml | 246 |
| methotrexate sodium oral tablet 2.5 mg.. | 246 |
| methyprednisolone oral tablet 16 mg, 32 | |
| mg, 4 mg, 8 mg | 246 |
| metronidazole intravenous solution 500 | |
| mg/100ml | 246 |
| miglustat..... | 127 |
| modafinil | 31 |
| moxifloxacin hcl in nacl intravenous | |
| solution 400 mg/250ml | 246 |
| mycophenolate mofetil oral capsule 250 mg | |
| | 246 |
| mycophenolate mofetil oral suspension | |
| reconstituted 200 mg/ml | 246 |

mycophenolate mofetil oral tablet 500 mg 246
mycophenolate sodium oral tablet delayed
release 180 mg, 360 mg 246

N

nafcillin sodium injection solution
reconstituted 1 gm, 2 gm..... 246
nafcillin sodium intravenous solution
reconstituted 10 gm..... 246
NAMZARIC 129
NATPARA..... 130
NERLYNX 131
NINLARO..... 132
nitisinone..... 133
NOXAFIL ORAL PACKET..... 154
NOXAFIL ORAL SUSPENSION..... 154
NUBEQA..... 134
NUCALA..... 135
NUEDEXTA..... 136
NUPLAZID ORAL CAPSULE..... 137
NUPLAZID ORAL TABLET 10 MG.... 137
NUTRILIPID INTRAVENOUS
EMULSION 20 %..... 246

O

octreotide acetate injection solution 100
mcg/ml, 1000 mcg/ml, 200 mcg/ml, 50
mcg/ml, 500 mcg/ml 138
ODOMZO 139
OFEV 140
OMNITROPE SUBCUTANEOUS
SOLUTION CARTRIDGE..... 77, 78
OMNITROPE SUBCUTANEOUS
SOLUTION RECONSTITUTED... 77, 78
ondansetron hcl oral solution 4 mg/5ml.. 246
ondansetron hcl oral tablet 4 mg, 8 mg... 246
ondansetron oral tablet dispersible 4 mg, 8
mg 246
ONUREG..... 141
OPSUMIT 142
ORFADIN ORAL CAPSULE 20 MG.... 133
ORGOVYX..... 143
ORKAMBI..... 144
OSPHENA 145
oxacillin sodium in dextrose intravenous
solution 1 gm/50ml, 2 gm/50ml..... 246

oxacillin sodium injection solution
reconstituted 1 gm, 2 gm..... 247
oxacillin sodium intravenous solution
reconstituted 10 gm..... 247
oxandrolone oral 146

P

PANRETIN..... 147
PANZYGA INTRAVENOUS SOLUTION
1 GM/10ML, 10 GM/100ML, 2.5
GM/25ML, 20 GM/200ML, 30
GM/300ML, 5 GM/50ML 247
paricalcitol oral capsule 1 mcg, 2 mcg, 4
mcg..... 247
PEGASYS SUBCUTANEOUS SOLUTION
180 MCG/ML 148
PEGASYS SUBCUTANEOUS SOLUTION
PREFILLED SYRINGE 148
PEMAZYRE..... 149
penicillin g potassium injection solution
reconstituted 20000000 unit..... 247
penicillin g sodium injection solution
reconstituted 5000000 unit..... 247
pentamidine isethionate inhalation solution
reconstituted 300 mg..... 247
pentamidine isethionate injection solution
reconstituted 300 mg..... 247
perphenazine oral tablet 4 mg, 8 mg..... 247
PIQRAY (200 MG DAILY DOSE)..... 150
PIQRAY (250 MG DAILY DOSE)..... 150
PIQRAY (300 MG DAILY DOSE)..... 150
pirfenidone oral tablet 151
PLASMA-LYTE 148 INTRAVENOUS
SOLUTION..... 247
PLASMA-LYTE A INTRAVENOUS
SOLUTION..... 247
POMALYST 152
posaconazole 153
potassium chloride in nacl intravenous
solution 20-0.45 meq/l-%, 20-0.9 meq/l-
%, 40-0.9 meq/l-% 247
potassium chloride intravenous solution 2
meq/ml, 2 meq/ml (20 ml), 20 meq/100ml
..... 247
potassium cl in dextrose 5% intravenous
solution 20 meq/l..... 247
prednisolone oral solution 15 mg/5ml 247

March 2023

H9306_23_DRS_006_OE_C

| | |
|---|-----|
| prednisolone sodium phosphate oral solution 10 mg/5ml, 20 mg/5ml, 25 mg/5ml, 6.7 (5 base) mg/5ml..... | 247 |
| prednisolone sodium phosphate oral tablet dispersible 10 mg, 15 mg, 30 mg..... | 247 |
| PREDNISONE INTENSOL ORAL CONCENTRATE 5 MG/ML..... | 247 |
| prednisone oral solution 5 mg/5ml | 247 |
| prednisone oral tablet 1 mg, 10 mg, 2.5 mg, 20 mg, 5 mg, 50 mg | 247 |
| prehevbrio intramuscular suspension 10 mcg/ml | 247 |
| PREMASOL INTRAVENOUS SOLUTION 10 % | 247 |
| PREVYMIS ORAL | 155 |
| PRIVIGEN INTRAVENOUS SOLUTION 20 GM/200ML | 247 |
| prochlorperazine maleate oral tablet 10 mg, 5 mg | 247 |
| PROGRAF ORAL PACKET 0.2 MG, 1 MG | 247 |
| PROLASTIN-C INTRAVENOUS SOLUTION RECONSTITUTED | 6 |
| PROMACTA | 156 |
| PROSOL INTRAVENOUS SOLUTION 20 % | 247 |
| PULMOZYME INHALATION SOLUTION 2.5 MG/2.5ML | 247 |
| Q | |
| QINLOCK | 157 |
| quinine sulfate oral..... | 158 |
| R | |
| RABAVERT INTRAMUSCULAR SUSPENSION RECONSTITUTED... .. | 247 |
| RAVICTI | 159 |
| RECOMBIVAX HB INJECTION SUSPENSION 10 MCG/ML, 40 MCG/ML, 5 MCG/0.5ML | 247 |
| RECOMBIVAX HB INJECTION SUSPENSION PREFILLED SYRINGE 10 MCG/ML, 5 MCG/0.5ML | 247 |
| REGRANEX | 160 |
| REPATHA | 161 |
| REPATHA PUSHTRONEX SYSTEM.. | 161 |
| REPATHA SURECLICK | 161 |

| | |
|--|-----|
| RETACRIT INJECTION SOLUTION 10000 UNIT/ML, 10000 UNIT/ML(1ML), 2000 UNIT/ML, 20000 UNIT/ML, 3000 UNIT/ML, 4000 UNIT/ML, 40000 UNIT/ML | 56 |
| RETEVMO | 162 |
| REZUROCK | 163 |
| riluzole | 164 |
| RINVOQ | 165 |
| ROZLYTREK | 166 |
| RUBRACA | 167 |
| RYDAPT | 168 |
| S | |
| sapropterin dihydrochloride oral packet . | 169 |
| sapropterin dihydrochloride oral tablet... | 169 |
| SCEMBLIX | 170 |
| SIGNIFOR | 171 |
| sildenafil citrate oral tablet 20 mg | 172 |
| sirolimus oral solution 1 mg/ml | 247 |
| sirolimus oral tablet 0.5 mg, 1 mg, 2 mg | 247 |
| SIRTURO | 173 |
| SKYRIZI PEN | 174 |
| SKYRIZI SUBCUTANEOUS | 174 |
| sofosbuvir-velpatasvir..... | 79 |
| SOLTAMOX | 175 |
| SOMAVERT | 176 |
| sorafenib tosylate | 177 |
| SPRYCEL | 178 |
| STELARA SUBCUTANEOUS SOLUTION 45 MG/0.5ML | 179 |
| STELARA SUBCUTANEOUS SOLUTION PREFILLED SYRINGE | 179 |
| STIVARGA | 180 |
| sunitinib malate..... | 181 |
| SUNOSI | 182 |
| SYMDEKO | 183 |
| SYMLINPEN 120 SUBCUTANEOUS SOLUTION PEN-INJECTOR..... | 184 |
| SYMLINPEN 60 SUBCUTANEOUS SOLUTION PEN-INJECTOR..... | 184 |
| SYNAREL | 185 |
| SYNRIBO | 186 |
| T | |
| TABLOID | 187 |
| TABRECTA | 188 |

| | |
|---|-----|
| tacrolimus oral capsule 0.5 mg, 1 mg, 5 mg | 247 |
| TAFINLAR..... | 189 |
| TAGRISSO..... | 190 |
| TAKHZYRO..... | 191 |
| TALZENNA | 192 |
| TASIGNA..... | 193 |
| TAVNEOS..... | 194 |
| tazarotene external cream | 195 |
| tazarotene external gel | 195 |
| TAZORAC EXTERNAL CREAM 0.05 % | 195 |
| TAZVERIK..... | 196 |
| TDVAX INTRAMUSCULAR SUSPENSION 2-2 LF/0.5ML | 247 |
| TEFLARO INTRAVENOUS SOLUTION RECONSTITUTED 400 MG, 600 MG | 247 |
| TEGSEDI..... | 197 |
| TENIVAC INTRAMUSCULAR INJECTABLE 5-2 LFU, 5-2 LFU (INJECTION) | 247 |
| TEPMETKO | 198 |
| teriparatide (recombinant)..... | 199 |
| tetrabenazine | 200 |
| THALOMID | 201 |
| TIBSOVO | 202 |
| tigecycline intravenous solution reconstituted 50 mg..... | 247 |
| TOBI PODHALER..... | 203 |
| tobramycin inhalation nebulization solution 300 mg/5ml | 247 |
| tobramycin sulfate injection solution 10 mg/ml, 80 mg/2ml..... | 247 |
| tolvaptan..... | 204 |
| toremifene citrate | 206 |
| TPN ELECTROLYTES INTRAVENOUS CONCENTRATE | 247 |
| TRAVASOL INTRAVENOUS SOLUTION 10 % | 247 |
| TRELSTAR MIXJECT | 207 |
| tretinoin external | 205 |
| TREXALL ORAL TABLET 10 MG, 15 MG, 5 MG, 7.5 MG | 247 |
| trientine hcl | 208 |
| TRIKAFTA..... | 209 |

| | |
|--|-----|
| TROPHAMINE INTRAVENOUS SOLUTION 10 % | 247 |
| TRUSELTIQ (100MG DAILY DOSE).. | 210 |
| TRUSELTIQ (125MG DAILY DOSE).. | 210 |
| TRUSELTIQ (50MG DAILY DOSE).... | 210 |
| TRUSELTIQ (75MG DAILY DOSE).... | 210 |
| TUKYSA | 211 |
| TURALIO ORAL CAPSULE 200 MG.. | 212 |
| TYMLOS | 213 |
| U | |
| UBRELVY..... | 214 |
| V | |
| VALCHLOR..... | 215 |
| VARUBI (180 MG DOSE) ORAL TABLET THERAPY PACK 2 X 90 MG | 247 |
| VENCLEXTA..... | 216 |
| VENCLEXTA STARTING PACK | 216 |
| VERQUVO | 217 |
| VERZENIO..... | 218 |
| vigabatrin | 219 |
| VIJOICE | 220 |
| VITRAKVI | 221 |
| VIZIMPRO | 222 |
| VONJO | 223 |
| voriconazole intravenous | 224 |
| voriconazole oral..... | 224 |
| VOSEVI..... | 79 |
| VOTRIENT..... | 225 |
| VYNDAMAX..... | 226 |
| W | |
| WELIREG..... | 227 |
| X | |
| XALKORI..... | 228 |
| XATMEP ORAL SOLUTION 2.5 MG/ML | 248 |
| XGEVA..... | 229 |
| XOLAIR | 230 |
| XOSPATA | 231 |
| XPOVIO (100 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 50 MG | 232 |
| XPOVIO (40 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 40 MG.. | 232 |

| | |
|--|-----|
| XPOVIO (40 MG TWICE WEEKLY) ORAL TABLET THERAPY PACK 40 MG | 232 |
| XPOVIO (60 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 60 MG.. | 232 |
| XPOVIO (60 MG TWICE WEEKLY)... | 232 |
| XPOVIO (80 MG ONCE WEEKLY) ORAL TABLET THERAPY PACK 40 MG.. | 232 |
| XPOVIO (80 MG TWICE WEEKLY)... | 232 |
| XTANDI | 233 |
| XURIDEN..... | 234 |
| XYREM | 235 |

| | |
|---------------------------|-----|
| XYWAV | 236 |
| Y | |
| YONSA..... | 237 |
| Z | |
| ZARXIO | 238 |
| ZEJULA..... | 239 |
| ZELBORAF | 240 |
| ZIEXTENZO | 241 |
| ZOLINZA | 242 |
| ZYDELIG | 243 |
| ZYKADIA ORAL TABLET | 244 |